

Changes of Pre Ejection Period and Left Ventricular Ejection Time during Head up Tilt

Ahlam Kadhim Al Hamdany^{1*}

¹Department of Physiology, College of Medicine, Babylon University, Iraq.

Author's contribution

The sole author designed, analyzed and interpreted and prepared the manuscript.

Article Information

DOI: 10.9734/CA/2015/12229

Editor(s):

(1) Francesco Pelliccia, Department of Heart and Great Vessels University La Sapienza, Rome, Italy.

(2) Anonymous

Reviewers:

(1) Andrea Borghini, Institute of Clinical Physiology- CNR, Pisa, Italy.

(2) Anonymous, Ospedale SS Annunziata di Cento (FE), Italy.

(3) Anonymous, Juntendo University, Japan.

(4) Anonymous, Sao Paulo University, Brazil.

Peer review History: <http://www.sciencedomain.org/review-history.php?iid=641&id=26&aid=5972>

Short Research Article

Received 24th June 2014
Accepted 19th August 2014
Published 6th September 2014

ABSTRACT

Objective of the Study: Evaluate changes in Pre ejection period (PEP) and left ventricular ejection time (LVET) during head up tilt (HUT).

Methods: Twenty healthy male subjects were involved in this study, with mean age 29.3±5 years, mean body mass index (BMI) 21.3±0.2Kg/m². Measurement of PEP, LVET of Doppler wave form of the aortic flow were done at supine, 30 and 60 degree HUT. Measurement of HR and BP were done at these positions of tilting. Comparison of changes of these variables at different degrees of HUT was done by paired T-Test.

Results: PEP values were significantly higher in 60 degree and 30 degree HUT than PEP values at supine position (p<0.0001). PEP values at 60 degree HUT were significantly higher than PEP values at 30 degree HUT (P=0.05).

LVET values were significantly lower at 60 degree and 30 degree HUT than values at supine position (p<0.001), and LVET values were significantly lower at 60 degree HUT than values at 30 degree HUT (p<0.001).

Conclusion: Key findings of PEP and LVET during HUT are progressive prolongation of PEP and shortening of LVET with increasing head up tilting.

Keywords: HUT; LVET; PEP.

*Corresponding author: E-mail: udayjanabia@yahoo.com;

ABBREVIATIONS

BMI-body mass index, BP-blood pressure, DBP-diastolic blood pressure, LVET-left ventricular ejection time, PEP-pre ejection period, SBP-systolic blood pressure, TTT-tilt table test.

1. INTRODUCTION

Tilt table test (TTT): TTT, over half a century old, has retained a central place in the investigation of syncope of unknown origin [1-7]. Since the differential diagnosis of syncope of unknown origin is widely spread, there have been many attempts to rationalize and improve the diagnostic procedure [8]. During HUT—or for that matter while standing—a person's cardiovascular system has to adjust itself in order to prevent a significant portion of the blood volume from pooling in the legs.

These adjustments consist of an increase in heart rate, and a constriction of the blood vessels in the legs. These cardiovascular adjustments occur very quickly, and there is no significant drop in the blood pressure [9]. The increase in heart rate of approximately 10-15 beats/min, an elevation of diastolic pressure of about 10mm Hg and little change in systolic pressure [10]. Many previous studies enlighten the changes that occurred in stroke volume, cardiac output, heart rate, and blood pressure during HUT. The aim of this study is to evaluate the changes that occur in LVET and PEP during different degree of tilting.

2. MATERIALS AND METHODS

2.1 Subjects

Twenty healthy male subjects were involved in this study, after having their signed consent and approval of the ethical committee at Kufa faculty of medicine, with mean age 29.3 ± 5 years, BMI $21.3 \pm 0.2 \text{ Kg/m}^2$. The subjects were thoroughly examined clinically to confirm the inclusion criteria (that only healthy subjects were included). Exclusion criteria included hypertension, diabetes mellitus, coronary heart disease and other cardiac problems, renal disease. None was on any medication.

2.2 Apparatus

2.2.1 Tilting table

A motorized tilt table. All TTT were performed in quite air conditioned room, especially equipped for the investigation [11].

2.2.2 Echocardiography equipment

All echocardiographic and Doppler studies were performed using two-dimensional (2D) Philips Sonos 7500 equipment with 2.5 MHZ transducer with tissue harmoni, incorporated ECG, and Doppler facilities for measurement of PEP and LVET.

2.3 Methods

To avoid any possible emotional excitement, reassurance of the subjects were done, of being safe and non invasive procedure.

Participants were examined in a quiet, temperature-controlled room, and were first supine positioned on belt secured tilting table, for at least 10-min to achieve a steady state. A steady state means that heart rate in consecutive minutes changes by less than 3 beats/min [12]. Pulse oximeter (portable battery oximeter-nonin-USA) was fixed on right index so as to digitally follow up the changes in arterial pulse to gain a steady state, and to record the HR. After reaching the steady state, we placed the transducer on the apex to get apical view and we used the continues wave Doppler for aorta. From aortic flow, PEP was estimated from Q-wave in ECG to the opening of aortic valve, estimation of LVET time was done between opening and closure of aortic valve [13]. Several measurements for PEP and LVET were made, then taking the mean for them. After that we raised the subject to 30 degree HUT and wait till reaching steady state by examining the pulseoximeter, then The same parameters were measured again. Returning the subject to supine position followed by raising the subject to 60 degree HUT and same previous measurements were done. Measurement of blood pressure were done at supine, 30 and 60 degree HUT.

2.4 Statistical Analysis

All values were expressed as mean \pm SD. Comparison between PEP, LVET HR, SBP, DBP and MBP, at supine position and different degrees of HUT were done by paired t-test. $p < 0.05$ was considered statistically significant and $P < 0.001$ was considered statistically highly significant.

3. RESULTS

All hemodynamic parameters were expressed as mean±SD, at different degrees of HUT, at Table 1.

PEP values were significantly higher in 60 degree and 30 degree HUT than PEP values at supine position ($p<0.0001$), and PEP values at 60 degree HUT were significantly higher than PEP values at 30 degree HUT ($P=0.05$).

LVET values were statistically lower at 60 degree and 30 degree HUT than values at supine position ($p<0.001$), and LVET values were significantly lower at 60 degree HUT than values at 30 degree HUT ($p<0.001$). HR values were significantly higher at 30 degree and 60degree HUT than values at supine position, and HR values at 60 degree HUT were significantly higher than HR values at 30 degree HUT ($P<0.001$).

There were no statistically significant differences between SBP values at supine and 30 and 60 degree HUT ($P>0.05$).

DBP and MBP values were significantly higher at 30 degree and 60 degree HUT than values at supine position ($p=0.05$, $p<0.001$) respectively. DBP and MBP values at 60 degree were significantly higher at 60 degree HUT than values at 30 degree HUT ($P=0.05$) Fig. 1.

4. DISCUSSION

The important findings of this study were the progressive reduction of LVET and prolongation of PEP with increasing tilting, which reflect a decline in central blood volume [14-18]. This reduction of central blood volume are due to pooling about 300-800ml of blood to the lower extremities by effect of gravity during HUT [19,20]. Prolongation of PEP and shortening of LVET were more pronounced at 60 degree HUT than 30 degree HUT, this could be explained that increasing tilting leads to more pooling of blood to the lower extremities and reduction of cardiac preload. This is in agreement with other studies like Chan et al. [21] who found that there was a significant increase in PEP at (20-30 degrees) HUT.

Table 1. Hemodynamic parameters at different degrees of HUT (values were expressed as mean±SD)

60 degree HUT	30 degree HUT	Supine	Parameters
90.20±20.5	82.3±16	61.35±14.17	PEP(ms)
240.6±19.1	267.18±20.11	307.5±26.8	LVET(ms)
83.64±9.14	73.11±8.98	69±8.78	HR(beat/min)
126.9±10.88	127.7±7.8	126±7.64	SBP(mmHg)
83.58±8.17	79.47±5.68	77.1±5.4	DBP(mmHg)
98±64±7.59	96.41±41	93±5.51	MBP(mmHg)




Fig. 1. Hemodynamic parameters during supine, 30 and 60 degree head up tilt

In 2008, Chan et al. [22] found that there was a significant decrease in LVET during HUT. Fucà G et al. [23] in 2011, assessed systolic ejection time as a hemodynamic marker of incipient bradycardiac vasovagal syncope, also found that ET significantly decreased throughout tilt testing.

Pooling of blood during HUT will cause stimulation of baroreceptors and increase sympathetic activity which will cause increase cardiac contractility, increase HR, and vasoconstriction [24-29]. A 60° HUT maximize passive orthostatic stress on the sympathetic system by blocking the influence of inferior limb musculoskeletal contractions that could increase venous return [30]. For that reason, maximum sympathetic activity are at 60 degree HUT, so HR and diastolic BP and MBP values were significantly higher at 60 degree HUT than values at supine and 30 degree HUT. There were no differences in SBP values at different degrees of tilting due to increased sympathetic activity that compensate for the reduction of cardiac output.

5. CONCLUSION

Key findings of PEP and LVET during HUT are progressive prolongation of PEP and shortening of LVET with increasing HUT.

CONSENT

All authors declare that 'written informed consent was obtained from the patient (or other approved parties) for publication of this case report and accompanying images.

ETHICAL APPROVAL

All authors hereby declare that all experiments have been examined and approved by the appropriate ethics committee and have therefore been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki.

ACKNOWLEDGEMENTS

Special thanks to Dr. Akeel AMH Zwain who support me and helped me during collection of data.

COMPETING INTERESTS

Author has declared that no competing interests exist.

REFERENCES

1. Low PA. Composite autonomic scoring scale for laboratory quantification of generalized autonomic failure. *Mayo Clin Proc.* 1993;68:748-52.
2. Schondorf R, Low PA. Idiopathic postural orthostatic tachycardia syndrome: An attenuated form of acute pandysautonomia? *Neurology.* 1993;43:132-7.
3. Kaufmann H. Neurally mediated syncope: Pathogenesis, diagnosis, and treatment. *Neurology.* 1995;45(suppl 5):12-8.
4. Benditt DG, Ferguson DW, Grubb BP, Kapoor WN, Kugler J, Lerman BB, et al. Tilt table testing for assessing syncope. *J Am Coll Cardiol.* 1996;28(1):263-75.
5. Grubb BP, Kosinski D. Current trends in etiology, diagnosis, and management of neurocardiogenic syncope. *Curr Opin Cardiol.* 1996;11:32-41
6. Linzer M, Yang EH, Estes NAM, Wang P, Vorperian VR, Kapoor WN. Diagnosing syncope. Part 1: Value of history, physical examination, and electrocardiography. *Ann Intern Med.* 1997;126(12):989-96.
7. Kapoor WN. Using a tilt table to evaluate syncope. *Am J Med Sci.* 1999;317(2):110-6.
8. Brignole M, Alboni P, Benditt D, Bergfeldt L, Blanc J, Bloch Thomsen PE, Van Dijk JG, Fitzpatrick A, Hohnloser S, Janousek J, Kapoor W, Kenny RA, Kulakowski J, Moya A, Raviele A, Sutton R, Theodorakis G, Weilling W. For the task force on syncope. European society of cardiology. Guidelines on management of syncope. *European Heart Jour.* 2002;22:1256-1306.
9. Richard NF. Tilt table testing. *About. com Health's Disease and Condition;* 2003.
10. Grubb BP, Kosinski D. Dysautonomic and reflex syncope syndromes. *Cardiol Clin.* 1997;15(2):257-68.
11. Baron-Esquivias G, Martinez-Rubio AM. Tilt table test: State of the Art. *Indian pacing Electrophysiol J.* 2003;3(4):239-252.
12. Hainsworth R, AL-Shamma YHH. Cardiovascular responses to stimulation of carotid baroreceptor in healthy subjects. *Clinical Science.* 1988;75:159-165.
13. Skinner JR, Roy RJ, Heads A, Hey EN, Hunter S. Estimation of pulmonary arterial pressure in the new born: Study of the repeatability of four Doppler

- echocardiographic techniques. *Pediatr Cardiol.* 1996;17:360-369.
14. Stafford RW, Harris WS, Weissler AM. Left ventricular systolic time intervals as indices of postural circulatory stress in man. *Circulation.* 1970;41:485-492.
 15. Geeraerts T, Albaladejo P, Declere AD, Duranteau J, Sales JP, Benhamou D. Decrease in left ventricular ejection time on digital arterial waveform during simulated hypovolemia in normal humans. *J Trauma.* 2004;56:845-849.
 16. Chan GS, Middleton PM, Celler BG, Wang L, Lovell NH. Automatic detection of left ventricular ejection time from a finger photoplethysmographic pulse oximetry waveform: Comparison with Doppler aortic measurement. *Physiol Meas.* 2007;28:439-452.
 17. Chan GS, Middleton PM, Celler BG, Wang L, Lovell NH. Change in pulse transit time and pre-ejection period during head-up tilt-induced progressive central hypovolaemia. *J Clin Monit Comput.* 2007;21:283-293.
 18. Chan GS, Middleton PM, Celler BG, Wang L, Lovell NH. Detecting change in left ventricular ejection time during head-up tilt-induced progressive central hypovolaemia using a finger photoplethysmographic pulse oximetry wave form. *J Trauma.* 2008;64:390-397
 19. Grubb BP, Kosinski D. Dysautonomic and reflex syncope syndromes. *Cardiol. Clin.* 1997;15(2):257-68.
 20. Grubb BP, Kimmels. Head-upright tilt table testing using a safe and easy way to assess neurocardiogenic syncope. *Postgrad. Med.* 1998;103(1):133-140.
 21. Chan GS, Middleton PM, Celler BG, Wang L, Lovell NH. Change in pulse transit time and pre-ejection period during head-up tilt-induced progressive central hypovolaemia. *J Clin Monit Comput.* 2007;21(5):283-93. Epub 2007 Aug 16.
 22. Chan GS, Middleton PM, Celler BG, Wang L, Lovell NH. Detecting change in left ventricular ejection time during head-up tilt-induced progressive central hypovolaemia using a finger photoplethysmographic pulse oximetry wave form. *J Trauma.* 2008;64:390-397.
 23. Fucà G, Dinelli M, Gianfranchi L, Bressan S, Corbucci G, Alboni P. Assessment of systolic ejection time as a hemodynamic marker of incipient bradycardic vasovagal syncope. A pilot study. *Pacing Clin Electrophysiol.* 2011;34(8):954-62.
 24. Hainsworth R, Al-Shamma YMH. Cardiovascular responses to upright tilting in healthy subjects. *Clinical Science.* 1988;74:17-22.
 25. Grubb BP, Kosinski D. Dysautonomic and reflex syncope syndromes. *Cardiol Clin.* 1997;15(2):257-68.
 26. Grubb BP, Kimmels. Head-upright tilt table testing using a safe and easy way to assess neurocardiogenic syncope. *Postgrad. Med.* 1998;103(1):133-140.
 27. Sung RY, Dunz D, Yu GE, Yamme, Fok TF. Cerebral blood flow during vasovagal syncope induce By active standing or head-up tilt archives of disease in children. 2000;82(2):154-158.
 28. Vijayalakshri P, Veliath S, Moha M. Effect of head-up tilt on cardiovascular responses in normal young volunteers *Indian J. Physiol Pharmacol.* 2001;44(4): 467-472.
 29. AL-Shamma YMH, AL-Khawaja SAM, AL-Abidy JMR. Effect of upright tilting on cardiovascular reflexes using echocardiographic method for estimating cardiac output. *Kufa Med J.* 2002;5(2):85-96.
 30. Lamarre-Cliche M, Cusson J. The fainting patient: Value of head-upright tilt table test in adult patients with orthostatic intolerance. *CMA J.* 2001;164(3):372-376.

© 2015 AL-Hamdany; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history.php?iid=641&id=26&aid=5972>