

Asian Journal of Case Reports in Surgery

Volume 7, Issue 2, Page 388-392, 2024; Article no.AJCRS.119036

# Enteric Duplication Cysts in a 15 Year Old Children: A Rare Case Report

# Anil Kumar Sharma <sup>a</sup>, Pradeep Kumar Verma <sup>a</sup>, Kamal Bansal <sup>a</sup>, Rohit Jain <sup>a</sup> and Ashok Kumar Rajpura <sup>a\*</sup>

<sup>a</sup> JLN Medical College, Ajmer, Rajasthan, India.

#### Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

**Open Peer Review History:** 

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: https://www.sdiarticle5.com/review-history/119036

Case Report

Received: 01/05/2024 Accepted: 02/07/2024 Published: 26/07/2024

# ABSTRACT

**Introduction:** Enteric duplication cysts are rare congenital malformations in children and can develop anywhere along the gastrointestinal tract from mouth to the rectum, the ileum is the most common site followed by oesophagus, jejunum, stomach, and duodenum.

**Presentation of Case:** A 15-year-old boy was admitted to our hospital with the complaint of cramping intermittent pain mainly in his right lower abdomen for 2 days. Ultrasonogram of the abdomen showed a hollow structure in the right iliac fossa with perforation. After initial resuscitation emergency exploratory laparotomy was performed which revealed Enteric duplication cysts.

**Discussion:** Once the diagnosis of Enteric duplication cysts has been made, treatment is necessary to improve the symptoms and to prevent serious complications, including intestine obstruction, intussusception, or hematochezia.

**Conclusion:** In conclusion, this case of an intestinal duplication cyst presented with an acute abdomen. Although such conditions are more common in children and extremely rare in adults, they have to be considered for a differential diagnosis of acute abdomen.

<sup>\*</sup>Corresponding author: E-mail: drashokrajpura@gmail.com;

*Cite as:* Sharma, Anil Kumar, Pradeep Kumar Verma, Kamal Bansal, Rohit Jain, and Ashok Kumar Rajpura. 2024. "Enteric Duplication Cysts in a 15 Year Old Children: A Rare Case Report". Asian Journal of Case Reports in Surgery 7 (2):388-92. https://journalajcrs.com/index.php/AJCRS/article/view/555.

Keywords: Enteric duplication cysts; gastrointestinal tract; ultrasonography.

# **1. INTRODUCTION**

"Enteric duplication cysts are rare congenital malformations in children and can develop anywhere along the gastrointestinal tract, from the mouth to the rectum. The ileum is the most common site (33%) followed by the oesophagus (20%), colon (13%), jejunum (10%), stomach (7%), and duodenum (5%)" [1,2]. "They are usually detected prenatally or in the first years of life. Enteric duplication cysts are considered a very rare condition, particularly in adults" [3]. "Enteric duplication cysts are typically located at the mesenteric boundary of the gastrointestinal tract" [4]. "Enteric duplication cysts are believed to occur between the 4th and 8th weeks of embryonic development" [5]. "Their aetiology is still unknown; several theories have been proposed to explain their pathophysiology, but no single hypothesis can justify all duplications, locations, and associated anomalies. Split notochord theory is often postulated" [6,7].

"The duplication cysts of the small intestine can be categorized into three types based on the anatomy of the cyst: saccular cysts (Type I: 79%), tubular cysts (Type 2: 21%), and small intramural cysts. The saccular type is the most common and usually does not communicate with the native gastrointestinal tract. Tubular cysts are more commonly found in the colon. Small intramural cysts often occur near or at the ileocecal valve and protrude into the enteric lumen" [8].

"Ultrasonography is the most used imaging method for the diagnosis of enteric duplication cysts. Magnetic resonance and computed tomography scans are less frequently used but can be helpful in cases of difficult surgical approach" [9,10]. "The histopathological examination of the specimen is confirmatory for ascertaining the intestinal origin of the cyst, showing epithelial lining inside and a layer of smooth muscle in the wall, resembling some part of the gastrointestinal tract" [11] (Fig. 5).

#### 2. PRESENTATION OF CASE

A 15-year-old boy was admitted to our hospital emergency ward with a complaint of cramping intermittent pain mainly in the right lower abdomen for 2 days. The pain was acute in onset and gradually progressive in nature. The pain initially started in the right iliac fossa and then spread all over the abdomen. The pain was aggravated by food intake and was relieved after taking medications. The pain was also associated with vomiting. The patient had two episodes of bilious vomiting. He was afebrile on touch.



Fig. 1. Ultrasonogram of the abdomen: A cystic formation with a parietal stratification of three alternating layers (gut sign)

The patient was a known case of Down syndrome diagnosed in childhood at around 6 years of age. There was no past history of any acute illness and also there was no history of any surgery in the past. On examination, his pulse rate was 100/min and BP was 110/70 mmHg in the right arm in a supine position. Most laboratory investigations were within normal limits except for elevated white blood counts. On palpation, the patient had generalized tenderness all over the abdomen along with guarding and rigidity. The skiagram of the abdomen showed free gas under both domes of the diaphragm. Ultrasonogram of the abdomen showed a hollow structure in the right iliac fossa

with perforation and A cystic formation with a parietal stratification of three alternating layers (gut sign). (Fig. 1) After initial resuscitation emergency exploratory laparotomy was performed which revealed a duplication cyst originating from the mesenteric side of the ileum, sharing a common muscularis layer with the adjacent ileum (Figs. 2-4).

Post-operative course was uneventful and the patient was discharged from the hospital on post-operative day 5. Histopathological examination confirmed the diagnosis of an enteric duplication cyst arising from the ileum.



Fig. 2. A duplication cyst originating from the mesenteric side of the bowel, sharing a common muscularis layer with the adjacent bowel



Fig. 3. Surgical specimens

Anil et al.; Asian J. Case Rep. Surg., vol. 7, no. 2, pp. 388-392, 2024; Article no.AJCRS.119036



Fig. 4. Resected material A tubular duplication cyst protruding into the enteric lumen



Fig. 5. Histopathological examination of specimen: intestinal origin of the cyst, showing epithelial lining inside and a layer of smooth muscle in the wall

### 3. DISCUSSION

"Once the diagnosis of Enteric duplication cysts has been made, treatment is necessary to improve the symptoms and to prevent serious complications, including intestine obstruction, intussusception, or hematochezia. The treatment approach for most Enteric duplication cysts is surgical resection. Resection can be performed through traditional open-access approaches, as well as thoracoscopic and laparoscopic approaches" [12]. Currently, minimally invasive surgery is becoming the elective approach, and most of the cysts can be resected successfully.

Ultrasonography is the method of choice to diagnose gastrointestinal Enteric duplication cysts. Although a double-wall sign in a cyst is the most typical for diagnosis of Enteric duplication cysts. The diagnosis is confirmed by histological examination.

#### 4. CONCLUSION

In conclusion, this case of an intestinal duplication cyst presented with an acute abdomen. Although such conditions are more common in children and extremely rare in adults, they have to be considered for a differential diagnosis of acute abdomen. Surgery is the treatment of choice in intestinal duplication cysts to avoid complications.

# DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative Al technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image

generators have been used during writing or editing of manuscripts.

# CONSENT

As per international standards or university standards, patient consent has been collected and preserved by the authors.

# ETHICAL APPROVAL

As per international standard guidelines written ethical approval has been collected and preserved by the authors.

# **COMPETING INTERESTS**

Authors have declared that no competing interests exist.

# REFERENCES

- 1. Hur J, Yoon CS, Kim MJ, Kim OH. Imaging features of gastrointestinal tract duplications in infants and children: from esophagus to rectum. Pediatr Radiol. 2007;37:691–699.
- C Tiwari, H Shah, M Waghmare, et al. Cysts of gastrointestinal origin in children: Varied presentation Pediatr Gastroenterol Hepatol Nutr. 2017;20:94-99.
- Berrocal T, Hidalgo P, Gutiérrez J, De Pablo L, Rodríguez-Lemos R. Imagen radiológica de las duplicaciones del tubo digestivo. Radiología. 2004;46:282–292.
- 4. MH Okur, MS Arslan, S Arslan, et al. Gastrointestinal tract duplications in children Eur Rev Med Pharmacol Sci.

2014;18:1507-1512. View Record in Scopus Google Scholar.

- Sangüesa Nebot C, Llorens Salvador R, Carazo Palacios E, Picó Aliaga S, Ibañez Pradas V. Enteric duplication cysts in children: Varied presentations, varied imaging findings. Insights into Imaging. 2018 Dec;9:1097-106.
- Qi BQ, Beasley SW, Williams AK. Evidence of a common pathogenesis for foregut duplications and esophageal atresia with tracheo-esophageal fistula. Anat Rec. 2001;264:93–100.
- D'Agostino V, Castaldo A, Catelli A, Pesce I, Genovese S, Coppola L, et al. An ileal duplication cyst case report: from diagnosis to treatment. Radiology Case Reports. 2021 Jul 1;16(7):1597-602.
- Wardell S, Vidican DE. Ileal duplication cyst causing massive bleeding in a child. J Clin Gastroenterol. 1990;12:681–4. Available:https://doi.org/10.1097/00004836 -199012000-00018
- Sharma S, Yadav AK, Mandal AK, Zaheer S, Yadav D, Samie A. Enteric duplication cysts in children: A clinicopathological dilemma. J Clin Diagn Res. 2015;9:8–11.
- 10. Macpherson RI. Gastrointestinal tract duplications: Clinical, pathologic, etiologic, and radiologic considerations. Radiographics. 1993;13:1063–1080.
- 11. LM Van Zitteren, M Ruppert, B Op de Beeck, et al. Infected enteric duplication cyst BMJ Case Rep. 2017;21:220-221.
- Zhang Z, Huang X, Chen Q, Li D, Zhou Q, Huang J, et al. Small intestine duplication cyst with recurrent hematochezia: A case report and literature review. BMC Gastroenterology. 2021 Dec;21:1-5.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history: The peer review history for this paper can be accessed here: https://www.sdiarticle5.com/review-history/119036