



Evolution of Induce and Physical Mutagen on Vegetative Traits of Papaya (*Carica papaya* L.)

Veersain ^{a*}, Arvind Kumar ^a, Satya Prakash ^a,
Vipin Kumar ^b, Mukesh Kumar ^c, Upendra Maurya ^a
and Mohit Kumara Charu Shekhawat ^d

^a Department of Fruit Science, Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut, India.

^b Department of Vegetable Science, Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut, India.

^c Department of Genetics and Plant breeding, Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut, India.

^d University Agriculture University, Kota, Rajasthan, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jeai/2024/v46i92816>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/122140>

Original Research Article

Received: 20/06/2024

Accepted: 23/08/2024

Published: 28/08/2024

ABSTRACT

The experiment was conducted on Horticultural Research Centre of Sardar Vallabhbhai Patel University of Agriculture and Technology, located in Meerut, Uttar Pradesh. The experiment was laid out in RBD design and data was recorded on different growth trait of papaya. Two variety of

*Corresponding author: E-mail: Veersainarya1@gmail.com;

Cite as: Veersain, Arvind Kumar, Satya Prakash, Vipin Kumar, Mukesh Kumar, Upendra Maurya, and Mohit Kumara Charu Shekhawat. 2024. "Evolution of Induce and Physical Mutagen on Vegetative Traits of Papaya (*Carica Papaya* L.)". *Journal of Experimental Agriculture International* 46 (9):174-85. <https://doi.org/10.9734/jeai/2024/v46i92816>.

papaya Arka surya and Coorg honey dew, the seed of papaya was treated with Gamma and EMS mutagen and sown in field for evolution. The treatment detail of the experiment was T1, 5 Gy, T2, 10 Gy, T3, 15 Gy, T4, 0.25% EMS, T5, 0.50% EMS, T6, 0.75% EMS, T7, 5 Gy+0.25% EMS, T8, 10 Gy+0.50% EMS, and T9, 15 Gy+0.75% EMS. Finally, on the basis of result recorded that in both year days taken to seed germination and stem girth(cm), 10GR better performance over all treatment, while in number of leaves at flowering and petiole length (cm), 5GR better performance over all treatment. Whereas, In EMS and (Gamma rays + EMS), 0.25% EMS and 5G+0.25%EMS perform better result for petiole length and number of leaves at flowering while for days taken to seed germination and stem girth, 0.50%EMS and 10G+0.50%EMS perform better result.

Keywords: Papaya; gamma rays; EMS; RBD.

1. INTRODUCTION

Papaya (*Carica papaya* L.), belonging to the Caricaceae family, is a key economic fruit crop cultivated across the tropical and subtropical regions globally. Originating from Mexico to Panama, this tropical fruit is known for its three sex forms: male, female, and hermaphrodite, making it a polygamous species. It has a relatively small diploid genome, consisting of 372 Mbp/1C and nine pairs of chromosomes [1] The fruit is highly nutritious, rich in vitamin A with 2020 IU per 100 grams is the 2nd highest after mango, and provides significant medicinal benefits such as diuretic and heart stimulant [2]. Papain, an enzyme extracted from unripe papaya fruits, is utilized in various industries, including meat tenderization, textile processing, and cosmetics manufacturing.

One of the major fruit crops grown in India, papaya (*Carica papaya* L.), is grown across the world's tropical and subtropical zones. One of the main techniques used in plant breeding to develop a variety of crops is mutant breeding. The process of selection benefits from the creation of variability. Crops like grapes have improved greatly as a result of spontaneous mutations. Gamma irradiation is one of the physical mutagenic agents that may be used to boost crossing over frequencies (Celik et al., 2017). Breeders have long noted that the use of physical mutagens or mutagens may induce variety. Papaya cultivation thrives in countries like India, the USA, Mexico, Peru, Brazil, Jamaica, Nigeria, China, Taiwan, Indonesia, the Philippines, and Thailand. In India, papaya is grown across an area of 0.14 million hectares, resulting in a production of 5.74 metric tons per hectare [3]. Mutation breeding has proven effective for enhancing papaya traits, employing both chemical and physical mutagens to introduce genetic variability [4]. An example of successful mutation breeding is the development

of 'Pusa Nanha', previously known as the 'mutant dwarf' variety, by ICAR-IARI in New Delhi.

Mutagenesis is a valuable technique in crop improvement, involving the alteration of the genetic information of an organism to generate variability that can be harnessed for breeding new traits, such as enhanced nutritional qualities, resistance to diseases, or improved agronomic characteristics. In the case of papaya (*Carica papaya* L.), a popular tropical fruit, both qualitative and quantitative traits are significant for its commercial value, consumer preference, and adaptability to different growing conditions [5]. Mutation breeding helpful for improvement of different qualitative and quantitative traits, in case of qualitative trait, such as fruit colour, shape, and the presence or absence of specific diseases resistance genes governed by one or a few genes, while in quantitative traits yield, fruit size, sweetness, and overall plant growth vigour and environmental stress tolerance controlled by multiple genes.

Mutation breeding is a method used to create genetic diversity through novel recombinations, utilizing both chemical and physical mutagens. Pusananha, an ultra-dwarf variety with a dioecious nature, was developed by ICAR-IARI in New Delhi through mutation breeding using gamma irradiation. This approach could also be applied to develop gynodioecious varieties, with the potential to produce a dwarf mutant that is tolerant or resistant to Papaya Ring Spot Virus (PRSV), while also having good yield, quality, and extended shelf life. The current study examines the impact of gamma irradiation on the growth and yield parameters of papaya [6,7].

2. MATERIALS AND METHODS

The experiment was carried out at the Horticultural Research Centre of Sardar Vallabhbhai Patel University of Agriculture and

Technology, located in Meerut, Uttar Pradesh. In the experiment, papaya from two cultivars was used, namely Arka Surya and Coorg Honeydew, which were treated with gamma rays at varying levels (5, 10, and 15 Gy). The seed was treated with Nuclear Research Laboratory, IARI, New Delhi, and different levels of EMS concentrations (0.25%, 0.50%, and 0.75%) were treated with a diluted solution of water to achieve the necessary concentrations. After treatment, the seed with the mutagenic solution was placed in petri dishes, and the seeds were treated for 2 hours, and then the seed was washed in running water. Twenty seeds were placed in each treatment during the first year, and in the second year, the best quality fruit seed from each treatment was sown in the experimental field. However, the treatment combination of the experiment was T1, 5 Gy, T2, 10 Gy, T3, 15 Gy, T4, 0.25% EMS, T5, 0.50% EMS, T6, 0.75% EMS, T7, 5 Gy+0.25% EMS, T8, 10 Gy+0.50% EMS, and T9, 15 Gy+0.75% EMS. The analysis of variance for each character was carried out for the randomized block design as suggested by Panse and Sukhatme, [8] and with the help of opstat, Sheoran et al. [9].

3. RESULTS AND DISCUSSION

3.1 Vegetative Traits

3.1.1 Days taken to seed germination

Days to first germination in papaya cv. Coorg Honey Dew and Arka Surya presented Table 1 and Fig. 1 showed that the significant difference in all treatment for days taken germination in 2022-23, in Gamma rays 10GR (16.67, 16.50) take minimum days for germination followed by 5GR (18.83, 18.00), and the maximum days taken for germination 15GR (19.33, 19.17), while the in EMS reported that 0.50%EMS (17.33, 16.17) take minimum days for seed germination over 0.25%EMS (17.50, 17.50) and the maximum days taken by 0.75%EMS (19.67, 18.33). Whereas in both (Gamma rays + EMS) recorded that 10G+0.50%EMS (19.50, 18.50) was taken as the minimum day for germination, followed by 5G+0.25%EMS (19.67, 18.67) and 15G+0.75%EMS (20.83, 20.50) were taken as the maximum days for germination, while in the case of the control taken as (18, 17.17) days for seed germination. However, in the case of all mutagens recorded, 10GR took a minimum of days for germination, similar finding reported by Pujra et al., [10] and Shailendra et al., [11].

During 2023-24 recorded that significant differences in all treatment for days taken for germination, Gamma rays 10GR (17.77, 17.30) take minimum days for germination followed by 5GR (18.01, 17.50), and the maximum days taken for germination 15GR (18.55, 18.48), while the in EMS reported that 0.50 EMS (18.00, 17.40) take minimum days for seed germination over 0.25% EMS (18.30, 18.17) and the maximum days taken by 0.75%EMS (18.65, 18.58). Whereas in both (Gamma rays + EMS) recorded that 10G+0.50%EMS (18.36, 18.15) was taken as the minimum day for germination, followed by 5G+0.25%EMS (19.15, 18.36) and 15G+0.75%EMS (19.51, 18.56) were taken as the maximum days for germination, while in the case of the control taken as (17.45, 17.62) days for seed germination. However, in the case of all mutagens recorded, 10GR took a minimum of days for germination, as presented in Table 1 and Fig. 1, result was conformed with smitha et al., [6,7] and Santosh et al., [12].


However, the average of both years reported that Gamma rays 10GR (17.22, 16.90) take minimum days for germination, followed by 5GR (18.42, 17.75), and the maximum days taken for germination 15GR (18.94, 18.82), while the EMS reported that 0.50%EMS (17.67, 16.79) take minimum days for seed germination over 0.25%EMS (17.90, 17.84) and the maximum days taken by 0.75%EMS (18.66, 18.46). Whereas in both (Gamma rays + EMS) recorded that 10G+0.50%EMS (18.93, 18.33) was taken as the minimum day for germination, followed by 5G+0.25%EMS (19.41, 18.51) and 15G+0.75%EMS (20.17, 19.53) were taken as the maximum days for germination, while in the case of the control taken as (17.72, 17.39) days for seed germination. However, in the case of all mutagens recorded, 10GR took a minimum of days for germination, as presented in Table 1 and Fig. 1, closed result conformed with Nishimwe et al., [5] and Jayshree et al., [13].

3.1.2 No. of leaves at flowering


Data presented Table-1 and Fig. 2 showed significant differences in all treatment during 2022-23 in papaya cv Coorg honey dew and Arka Surya, in gamma rays 5GR (34.50, 29.33) had maximum number of leaves followed by 10GR (31.33, 26.17) and minimum number of leaves at 15GR (30.67, 25.00), while the EMS reported that 0.25%EMS (28.50, 23.33) had maximum number of leaves followed by 0.50%EMS (27.83, 22.17) and minimum number

of leaves observed at 0.75%EMS (26.67, 21.67), whereas in both (Gamma rays + EMS) recorded that 5G+0.25%EMS (25.67, 20.33) had maximum number of leaves over 10G+0.50%EMS (24.83, 19.17) and minimum number of leaves observed in 15G+0.75%EMS

(21.50, 16.00), while in the case of the control taken as (20.83, 15.17) number of leaves. However, in the case of all mutagens recorded, 5GR had maximum number of leaves at flowering, results conform with smitha et al., [6,7] and Santhosh et al., (2010).



(A)



(b)

Fig. 1. Effect of gama rays, EMS and both of them (Gamma+EMS) on days taken to seed germination percentage of papaya cv. coorg honey dew and arka surya


Table 1. Effect of gama rays, EMS and both of them (Gamma+EMS) on days taken to seed germination percentage and number of leaves at first flowering of papaya cv. coorg honey dew and arka surya

		Days taken to germination						Number of leaves at first flowering					
		Coorg honey dew			Arka Surya			Coorg honey dew			Arka Surya		
		2022-23	2023-24	Average mean	2022-23	2023-24	Average Mean	2022-23	2023-24	Average mean	2022-23	2023-24	Average mean
Treatment		Y ₁	Y ₂		Y ₁	Y ₂		Y ₁	Y ₂		Y ₁	Y ₂	
T1	5 GR	18.83	18.01	18.42	18.00	17.50	17.75	34.50	36.83	35.67	29.33	31.50	30.42
T2	10 GR	16.67	17.77	17.22	16.50	17.30	16.90	31.33	34.67	33.00	26.17	28.67	27.42
T3	15 GR	19.33	18.55	18.94	19.17	18.48	18.82	30.67	33.00	31.83	25.00	27.67	26.33
T4	0.25 EMS	17.50	18.30	17.90	17.50	18.17	17.84	28.50	31.83	30.17	23.33	25.50	24.42
T5	0.50EMS	17.33	18.00	17.67	16.17	17.40	16.79	27.83	29.67	28.75	22.17	24.67	23.42
T6	0.75 EMS	18.67	18.65	18.66	18.33	18.58	18.46	26.67	29.50	28.08	21.67	24.33	23.00
T7	5 G+ 0.25 E	19.67	19.15	19.41	18.67	18.36	18.51	25.67	27.83	26.75	20.33	22.50	21.42
T8	10 G+ 0.50 E	19.50	18.36	18.93	18.50	18.15	18.33	24.83	26.33	25.58	19.17	21.67	20.42
T9	15 G+ 0.75 E	20.83	19.51	20.17	20.50	18.56	19.53	21.50	23.67	22.58	16.00	18.83	17.42
Control	Control	18.00	17.45	17.72	17.17	17.62	17.39	20.83	22.83	21.83	15.17	17.67	16.42
SE(m±)		0.58	0.31		0.53	0.22		0.85	0.87		0.69	0.72	
C.D. at 5%		1.73	0.91		1.58	0.64		2.55	2.61		2.06	2.15	


Table 2. Effect of gama rays, EMS and both of them (Gamma+EMS) on stem girth (cm) and petiole length (cm) of papaya cv. coorg honey dew and arka surya

		Stem girth (cm)						Petiole length (cm)					
		Coorg honey dew			Arka Surya			Coorg honey dew			Arka Surya		
		2022-23	2023-24	Average mean	2022-23	2023-24	Average mean	2022-23	2023-24	Average mean	2022-23	2023-24	Average mean
Treatment		Y ₁	Y ₂		Y ₁	Y ₂		Y ₁	Y ₂		Y ₁	Y ₂	
T1	5 GR	38.18	40.23	39.21	40.23	42.62	41.43	40.32	39.88	40.10	36.58	37.05	36.82
T2	10 GR	39.43	41.32	40.38	41.32	45.67	43.49	38.38	37.67	38.03	33.73	35.22	34.48
T3	15 GR	36.38	38.13	37.26	38.13	40.12	39.13	38.23	36.88	37.56	33.52	35.07	34.29
T4	0.25 EMS	36.33	38.32	37.33	38.58	42.90	40.74	36.43	35.87	36.15	31.78	33.18	32.48
T5	0.50EMS	37.13	38.58	37.86	38.32	43.37	40.84	35.88	34.85	35.37	31.23	32.72	31.98
T6	0.75 EMS	35.42	37.13	36.28	37.13	41.63	39.38	34.95	33.93	34.44	30.57	32.07	31.32

		Stem girth (cm)						Petiole length (cm)					
		Coorg honey dew			Arka Surya			Coorg honey dew			Arka Surya		
		2022-23	2023-24	Average mean	2022-23	2023-24	Average mean	2022-23	2023-24	Average mean	2022-23	2023-24	Average mean
Treatment		Y ₁	Y ₂		Y ₁	Y ₂		Y ₁	Y ₂		Y ₁	Y ₂	
T7	5 G+ 0.25 E	30.20	32.23	31.22	32.23	34.85	33.54	34.18	32.83	33.51	29.53	31.02	30.28
T8	10 G+ 0.50 E	32.83	34.62	33.73	34.62	39.06	36.84	32.45	31.13	31.79	27.70	28.67	28.18
T9	15 G+ 0.75 E	29.03	30.68	29.86	30.68	33.77	32.23	31.68	30.50	31.09	27.03	27.82	27.43
Control	Control	27.93	30.07	29.00	30.07	32.02	31.04	42.43	41.13	41.78	37.17	36.73	36.95
SE(m±)		1.07	1.21		1.19	1.15		1.22	1.04		1.46	3.05	
C.D. at 5%		3.20	3.61		3.58	3.45		3.67	3.10		4.37	1.02	



(A)




(B)


Fig. 2. Effect of gama rays, EMS and both of them (Gamma+EMS) on number of leaves at flowering of papaya cv. coorg honey dew and Arka Surya

Data showing Table-1 and Fig. 2 exposed significant differences in all treatment during 2023-24 in papaya cv. Coorg honey dew and Arka Surya, in gamma rays 0.5GR (36.63, 31.50) had maximum number of leaves followed by 10GR (34.67, 28.67) and 15GR (33.00, 27.67) had minimum number of leaves at flowering, while the EMS reported that 0.25%EMS (31.83, 25.50) had maximum number of leaves followed by 0.50% EMS (29.67, 24.67) and minimum number of leaves observed 0.75% EMS (29.50,

24.33), whereas in both (Gamma rays + EMS) recorded that 5G+0.25%EMS (27.83, 22.50) had maximum number of leaves over 10G+0.50%EMS (26.33, 21.67) and 15G+0.75%EMS (23.67, 18.33) had minimum number of leaves, while in the case of the control taken as (22.83, 17.67) number of leaves at flowering. However, in the case of all mutagens recorded, 5GR had maximum of number of leaves at flowering, finding similar to Sing et al., [2] and Kumar et al., [14].



(A)



(B)

Fig. 3. Effect of gama rays, EMS and both of them (Gamma+EMS) on stem girth of papaya cv. coorg honey dew and arka surya

Number of leaves at flowering present Table 1 and Fig. 2 exposed significant differences in all treatment in both of year average, in gamma rays 0.5GR (35.67, 30.42) had maximum number of leaves at flowering followed by 10GR (33.00, 27.2) and 15GR (31.83, 26.33) minimum number of leaves at flowering, while the EMS reported that 0.25% EMS (30.17, 24.42) had maximum number of leaves at flowering followed by 0.50%EMS (28.75, 23.42) and minimum number of leaves observed 0.75%EMS (28.08, 23.00), whereas in both (Gamma rays + EMS) recorded that 5G+0.25%EMS (26.75, 21.42) had maximum number of leaves over


10G+0.50%EMS (25.58, 20.42) and 15G+0.75%EMS (22.58, 17.42) had minimum number of leaves at flowering, while in the case of the control taken as (21.83, 16.42) had number of leaves at flowering. However, in the case of all mutagens recorded, 5GR had maximum number of leaves at flowering, closed result with Jayshree et al., [13] and Kumar et al., [14].

3.1.3 Stem girth (cm)


Data presented in Table-1 and Fig. 3 exhibited significant differences in all treatment during

2022-23 on papaya cv Coorg honey dew and Arka Surya, in gamma rays, 10GR (39.43, 44.77) had maximum stem girth, followed by 5GR (38.18, 41.72), and the minimum stem girth were 15GR (36.38, 40.82), while the in EMS reported that 0.50% EMS (37.13, 41.97) had maximum stem girth over 0.25% EMS (36.33, 41.67) and the minimum stem girth observed by 0.75%EMS (35.42, 40.13). Whereas in both (Gamma rays + EMS) recorded that 10G+0.50%EMS (32.83,

38.17) had maximum stem girth, followed by 5G+0.25%EMS (30.20, 32.63) and 15G+0.75%EMS (29.03, 30.63) were taken as the minimum stem girth, while in the case of the control taken as (27.93, 29.45) had stem girth. However, in the case of all mutagens recorded, 10GR had the maximum stem girth, result conformed with Singh et al., [2] and Santosh et al., [12].



(A)



(B)

Fig. 4. Effect of gama rays, EMS and both of them (Gamma+EMS) on petiole length (cm) of papaya cv. coorg honey dew and arka surya

The data showed in Table 1 and Fig. 3 revealed significant differences in all treatment during 2023-24 in two papaya cv. Coorg honey dew and Arka Surya, gamma rays 10GR (41.32, 45.67) had maximum stem girth, followed by 5GR (40.23, 42.62), and the minimum stem girth were 15GR (38.13, 40.12), while the in EMS reported that 0.50%EMS (38.58, 43.37) had maximum stem girth over 0.25%EMS (38.32, 42.90) and the minimum stem girth observed by 0.75%EMS (37.13, 41.63). Whereas in both (Gamma rays + EMS) recorded that 10G+0.50%EMS (34.62, 39.06) had maximum stem girth, followed by 5G+0.25%EMS (32.23, 34.85) and 15G+0.75%EMS (30.68, 33.77) were taken as the minimum stem girth, while in the case of the control taken as (30.07, 32.02) had stem girth. However, in the case of all mutagens recorded, 10GR had the maximum stem girth, result similar with Kumar *et al.*, [14] and Pujari *et al.*, [10]

In average of both years reported that Gamma ray's gamma rays 10GR (40.38, 43.49) had maximum stem girth, followed by 5GR (39.21, 41.43), and the minimum stem girth were 15GR (37.26, 39.13), while the in EMS reported that 0.50% EMS (37.86, 40.84) had maximum stem girth over 0.25% EMS (37.33, 40.74) and the minimum stem girth observed by 0.75%EMS (36.28, 39.38). Whereas in both (Gamma rays + EMS) recorded that 10G+0.50%EMS (33.73, 36.84) had maximum stem girth, followed by 5G+0.25%EMS (31.22, 33.54) and 15G+0.75%EMS (29.86, 32.23) were taken as the minimum stem girth, while in the case of the control taken as (29.00, 31.04) had stem girth. However, in the case of all mutagens recorded, 10GR had the maximum stem girth, as presented in Table 2 and Fig. 3, result conformed with Shailendra *et al.*, [11] and Kumar *et al.*, [13].

3.1.4 Petiole length (cm)

Data existing in Table- 2 and Fig. 4 exhibited significant differences in all treatment during 2022-23 on papaya cv Coorg honey dew and Arka Surya, in gamma rays, 5GR (40.32, 36.58) had maximum petiole length, followed by 10GR (38.38, 33.73), and the minimum petiole length were 15GR (38.23, 33.52), while the in EMS reported that 0.25%EMS (34.95, 31.78) had maximum petiole length over 0.50%EMS (35.88, 31.23) and the minimum petiole length observed by 0.75%EMS (34.95, 30.57). Whereas in both (Gamma rays + EMS) recorded that

5G+0.25%EMS (34.18, 29.53) had maximum petiole length, followed by 10G+0.50%EMS (32.45, 27.70) and 15G+0.75%EMS (31.68, 27.03) were taken as the minimum petiole length, while in the case of the control taken as (42.43, 37.17) had petiole length. However, in the case of all mutagens recorded, 5GR had the maximum petiole length, result conformed with Singh *et al.*, [2].

The data showed in Table 2 and Fig. 4 revealed significant differences in all treatment during 2023-24 in two papaya cv. Coorg honey dew and Arka Surya, in gamma rays, 5GR (39.88, 37.05) had maximum petiole length, followed by 10GR (37.67, 35.22), and the minimum petiole length were 15GR (36.88, 35.07), while the in EMS reported that 0.25%EMS (35.87, 33.18) had maximum petiole length over 0.50%EMS (34.85, 32.72) and the minimum petiole length observed by 0.75%EMS (33.93, 32.07). Whereas in both (Gamma rays + EMS) recorded that 5G+0.25%EMS (32.83, 31.02) had maximum petiole length, followed by 10G+0.50%EMS (31.13, 28.67) and 15G+0.75%EMS (30.50, 27.82) were taken as the minimum petiole length, while in the case of the control taken as (41.13, 36.73) had petiole length. However, in the case of all mutagens recorded, 5GR had the maximum petiole length, finding conform with Kumar *et al.*, [13] and Bakry *et al.*, [14-17].

In average of both years reported that in gamma rays, 5GR (40.10, 36.82) had maximum petiole length, followed by 10GR (38.03, 34.48), and the minimum petiole length were 15GR (37.56, 34.29), while the in EMS reported that 0.25%EMS (36.15, 32.48) had maximum petiole length over 0.50%EMS (35.37, 31.98) and the minimum petiole length observed by 0.75% EMS (34.44, 31.32). Whereas in both (Gamma rays + EMS) recorded that 5G+0.25%EMS (33.51, 30.28) had maximum petiole length, followed by 10G+0.50%EMS (31.79, 28.18) and 15G+0.75%EMS (31.09, 27.43) were taken as the minimum petiole length, while in the case of the control taken as (41.78, 36.95) had petiole length. However, in the case of all mutagens recorded, 5GR had the maximum petiole length, as presented in Table 2 and Fig. 4, closed result with Kumar *et al.*, [14] and Singh *et al.*, [2].

4. CONCLUSION

Finally, on the basis of result concluded that in both year days taken to seed germination and

stem girth(cm), 10GR better performance over all treatment, while in number of leaves at flowering and petiole length (cm), 5GR better performance over all treatment. Whereas, In EMS and (Gamma rays + EMS), 0.25% EMS and 5G+0.25%EMS perform better result for petiole length and number of leaves at flowering while for days taken to seed germination and stem girth, 0.50%EMS and 10G+0.50%EMS perform better result.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Vincent L, Anushma PL, Vasugi C, Rekha A, Shiva B. Genetic resources of tropical fruits. Conservation and utilization of horticultural genetic resources. 2019;79-116.
- Singh, SV, Singh, DB, Yadav, M, Roshan, RK, & Pebam, N. Effect of EMS on germination, growth and sensitivity of papaya (*Carica papaya* L.) cv. Farm Selection-1. In II International Symposium on Papaya 851 2008;851:113-116.
- Anonymous, Agricultural statistics at glance. Directorate of economics and statistics, DAC & FW, Govt of India. 2022;78.
- Bado S, Yamba NG, Sesay JV, Laimer M, Forster BP. Plant mutation breeding for the improvement of vegetatively propagated crops: successes and challenges. CABI Reviews. 2018;3(2017):1-21.
- Nishimwe G. Characterization of morphological and quality characteristics of new papaya (*Carica papaya* L) Hybrids Developed at JKUAT (Doctoral dissertation, JKUAT-AGRICULTURE) 2019.
- Smitha S, Hanur VS, Shyamamma S. Field evaluation of gamma irradiated M1 population of papaya (*Carica papaya* L.) cv. arka prabhath, Mysore J. Agric. Sci., 2022;56(4):1-15.
- Smitha S, Hanur VS. Effect of Gamma Radiation for Improving Morphological Parameters in Papaya (*Carica papaya* L.) cv. Arka Prabhath in M1 Generation. Res. Jr. of Agril. Sci. 2022;13(4): 1235–1243.
- Panse, VG. and Sukhatme, PV. Statistical Methods for Agricultural Workers. Indian Council of Agricultural Research Publication, 1985;87-89.
- Sheoran, OP, Tonk, DS, Kaushik, LS, Hasija, RC and Pannu, RS. Statistics and Computer Applications by DS. Hooda & RC. Hasija Department of Mathematics Statistics, CCS HAU, Hisar 1998;139-143.
- Pujar DU, Vasugi C, Vageeshbabu HS, Honnabyraiah MK, Adiga D, Jayappa J. Evaluation of mutant progenies for improved morphological, fruit and yield traits. Journal of Pharmacognosy and Phytochemistry. 2019;8(3):2324-34.
- Shailendra R, Singh V, Singh DB, Murlee Y, Roshan RK, Nongallei P. Effect of E.M.S on germination, growth and sensitivity of Papaya (*Carica papaya* L.) cv. Farm Selection-1. Pro. Second international symposium. Acta Hort. 2010;851:113-116.
- Santosh LC, Dinesh MR, Rekha A. Mutagenic studies in papaya (*Carica papaya* L.), Acta Hort. 2010;851:109-112.
- Jayashree, MJ, Manamohan, M, and Hanur, VS. Effect of gamma irradiation on germination and survival of seedlings in papaya cv. Arka Prabhath. Mysore J. Agric. Sci., 2022;56(2):161-165.
- Kumar Mahesh, Kumar Mukesh, Prakash Satya, Gautam DK, Rao Sanjeev. Effect of seed treatment by ethyl methane sulphonate on growth, flowering and yield of papaya cv. Pusa Dwarf. Journal of Hill Agriculture 2016;7(1):64-68.
- Bakry KH. AE and Faten HM. Ismaeil. Chemicals ib. Presowing treatments of papaya seeds as influenced by some chemicals and irradiation on germination, growth, flowering, sex expression and fruit quality. Second international conference horticulture science. 2002;10-12.
- Çelik Ö, Atak Ç. Applications of ionizing radiation in mutation breeding. New

- insights on gamma rays. 2017;24(6):111-32.
17. Mahadevamma M, Dinesh MR, Kumari VR, Shankarappa TH. Evaluation of induced variability in papaya (*Carica papaya* L.) by physical mutagenesis. CIBTech J Biotech. 2012;1(1): 66-71.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/122140>