



Response of Chitosan Salicylic Acid Nanoformulations on Growth, Yield and Quality Attributes of Strawberry cv. Winter Dawn

Mahendra Kumar Gora ^{a*}, Virendra Singh ^a,
Vinod Saharan ^b, Rakesh Kumar Yadav ^c
and Surbhi Prithiani ^a

^a Department of Horticulture, Rajasthan College of Agriculture, MPUAT, Udaipur, Rajasthan-313001, India.

^b Department of Molecular Biology and Biotechnology, Rajasthan College of Agriculture, MPUAT, Udaipur, Rajasthan-313001, India.

^c Department of Horticulture, College of Agriculture, Ummadganj, Kota, AU, Kota-325001, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jsrr/2024/v30i102486>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/124686>

Original Research Article

Received: 02/08/2024

Accepted: 04/10/2024

Published: 08/10/2024

ABSTRACT

A field experiment was conducted during 2022-23 at horticulture farm, Rajasthan College of Agriculture, MPUAT, Udaipur, Rajasthan, India. The experiment was laid out in a Randomized Block Design with three replications. In this experiment, data showed that growth, yield and quality attributes of strawberry were significantly enhanced by Chitosan salicylic acid nanoformulation (CS-

*Corresponding author: E-mail: mahendra.k.gora@gmail.com;

Cite as: Gora, Mahendra Kumar, Virendra Singh, Vinod Saharan, Rakesh Kumar Yadav, and Surbhi Prithiani. 2024. "Response of Chitosan Salicylic Acid Nanoformulations on Growth, Yield and Quality Attributes of Strawberry Cv. Winter Dawn". *Journal of Scientific Research and Reports* 30 (10):606-10. <https://doi.org/10.9734/jsrr/2024/v30i102486>.

SA NF) @ 800-1200 ppm as compare to salicylic acid (SA) @ 100-300 ppm, bulk chitosan (BCH) @ 100-1200 ppm. According to the data the maximum values of growth attributes viz. plant height (25.92 cm), number of branches (11.84) and yield attributes viz. number of fruits per plant (28.68), fruit weight (20.02 g), yield per plant (659.65 g), estimated yield per hectare (366.47 qt.) and quality attributes viz. TSS (7.54 °Brix), ascorbic acid (62.91 mg/100g) and total sugar (7.30 %) were found maximum in CS-SA NF 800 ppm.

Keywords: Chitosan; CS-SA NF; salicylic acid; BCH; strawberry.

1. INTRODUCTION

The cultivated strawberry (*Fragaria X ananassa* Duch.) belongs to the family Rosaceae with chromosome number, $2n = 56$. It is one of the delicious and nutritious among soft fruits with 98 percent edible portion. Strawberry plants are stoloniferous herbs and spread via stolons or runners. Strawberry plant has a short thickened stem called “crown” and shallow root system. The leaves are trifoliate and arise from the crown. Fruit is a small berry developed from one ovule with multiple achenes called true fruit which is non-climacteric. Strawberry occupies a unique place among cultivated berry fruits, as its fruits are attractive, luscious, tasty nutritious with pleasant aroma and delicate flavor. Strawberry contains anticancer compound i.e., ellagic acid and vitamin- C [1]. Strawberry can be grown under sub-tropical climate and even at high altitude of tropical climate. It can be grown wide range of soil from heavy clays to light sand and gravels. However, strawberry plant grows well in sandy loam soil with pH 5.5-6.5. Strawberry requires an optimum day temperature of 22 °C and night temperature of 7 °C to 13 °C for maximum growth and development. In India, it is grown commercially in the states of Maharashtra, Punjab, Haryana, Karnataka, Madhya Pradesh, Jammu & Kashmir, Himachal Pradesh and Uttarakhand [2]. In India strawberry is cultivated in 1000 hectare with a production of 8000 MT [3]. Flower induction in strawberry is controlled primarily by the interaction of photoperiod temperature and genotype. An individual flower typically has 10 green sepals five white petals 20 to 30 stamens and 60 to 600 pistils. The greatest number of pistils occurs on the primary flower and decreases successively down the inflorescence. Each pistil contains a single ovary that develops into an achene [4]. The achenes (the true fruits of strawberry) together with receptacle form an aggregate fruit which is referred to as a berry (the eterio of achenes, which is an aggregate of achene fruitlets developed from a single flower) but it is not a true berry in the botanical sense [5].

Chitosan, a biopolymer derived from biowaste of sea food industry and soluble in organic acids. It is considered environment-friendly for agricultural uses as it is easily degraded in the environment and nontoxic to humans. Chitosan and its derivatives have been reported to elicit natural defense responses in plants, and have been used as a natural compound to control pre- and post-harvest pathogenic diseases [6]. Salicylic acid (SA) or orthohydroxyl benzoic acid and related compounds belong to a diverse group of plant phenolics. SA is involved in the induction of plant defence mechanisms under different biotic and abiotic stresses. It has been reported that the exogenous application of SA increased the photosynthetic rate, exchange of gas, and accumulation of osmotic regulators (sugars, proline, and glycine betaine) and it enhanced the mineral and water relations of the plants, therefore leading to better plant growth [7].

2. MATERIALS AND METHODS

Data for growth parameters viz. plant height (cm) and No. of branches plant⁻¹ were recorded by five randomly selected plants from each plot at 60 and 90 days after transplanting. Fruit weight (g), No. of fruits plant⁻¹, total yield plant⁻¹ (g), estimated yield ha⁻¹ (kg) were calculated by five randomly selected plants from each plot at harvest. The TSS of ripe fruit juice was determined with the help of a hand refractometer (0-32 °B) by putting a few drops of juice on the prism. The refractometer calibrated with distilled water before use. The total soluble solid was expressed in °Brix. The percent total sugars present in strawberry fruits were determined by Lane and Eynon method, [8].

3. RESULTS

The plant height was significantly increased by foliar spray of CS-SA NFs 800 ppm (25.92 cm at 90 DAT) followed by CS-SA NFs 1200 ppm (25.77 cm) and BCH 1200 ppm (25.73 cm) as compared to control (16.64 cm). The number of branches was also significantly increased by CS-

SA NFs 800 ppm (11.84 at 90 DAT) followed by CS-SA NFs 1200 ppm (10.45 at 90 DAT) and BCH 1200 ppm (10.26 at 90 DAT) as compared to control (7.55 at 90 DAT) and other treatments (Table 1).

Number of fruits per plant was significantly increased by foliar spray of CS-SA NFs 800 ppm and CS-SA NFs 1200 ppm (28.68 and 28.53 fruits respectively) as compared to control (18.17

fruits). Fruit weight was also maximum in foliar spray of CS-SA NFs 800 ppm (20.02 g) followed by CS-SA NFs 1200 ppm (17.88 g) as compared to control (14.84 g) and other treatments. Moreover, total yield per plant (659.65 g), estimated yield per hectare (366.47 qt.) were significantly higher in 800 ppm CS-SA NFs followed by 1200 ppm CS-SA NFs (595.21 g and 330.67 qt.) as compared to control (324.42 g and 180.23 qt.) (Table 2).

Table 1. Effect of chitosan-salicylic acid NFs on growth parameters of strawberry cv. winter dawn

Treatment	Plant height At 90 DAT (cm)	Number of branches at 90 DAT	Number of fruits per plant
Control (water)	16.64±0.35 ^f	7.55±0.16 ^e	18.17±0.32 ^d
Salicylic acid (SA)			
SA (100 ppm)	19.29±0.01 ^e	8.36±0.00 ^{de}	19.02±0.01 ^d
SA (200 ppm)	21.17±0.56 ^{de}	8.75±0.23 ^d	22.47±0.52 ^c
SA (300 ppm)	21.93±0.09 ^{cd}	8.98±0.03 ^{cd}	22.57±0.08 ^c
Bulk chitosan			
BCH (100 ppm)	21.31±0.09 ^{de}	8.94±0.03 ^d	22.76±0.33 ^c
BCH (400 ppm)	23.42±0.07 ^{bcd}	9.04±0.02 ^{cd}	22.81±0.17 ^c
BCH (800 ppm)	23.86±0.72 ^{abc}	10.35±0.17 ^b	23.12±0.06 ^b ^c
BCH (1200 ppm)	25.73±0.44 ^{ab}	10.26±0.28 ^b	23.56±0.15 ^b ^c
CS-SA NFs			
CS-SA NF (100 ppm)	22.85±0.17 ^{cd}	8.51±0.06 ^{de}	23.12±0.06 ^b ^c
CS-SA NF (400 ppm)	23.97±0.78 ^{abc}	9.96±0.32 ^{bc}	24.93±0.71 ^b
CS-SA NF (800 ppm)	25.92±0.77 ^a	11.84±0.27 ^a	28.68±0.60 ^a
CS-SA NF (1200 ppm)	25.77±0.60 ^{ab}	10.45±0.24 ^b	28.53±0.58 ^a

The growth parameters were recorded at 90 days after transplanting (DAT). Each value is mean of triplicate and each replicate consisted of 5 plants. 1st foliar was done at 30 days and 2nd at 60 days after transplanting. Mean ± SE followed by same letter is not significantly different at $p = 0.05$ as determined by Tukey- Kramer HSD. BCH represents bulk chitosan dissolved in 0.5% acetic acid

Table 2. Effect of chitosan-salicylic acid NFs on yield attributes of strawberry cv. winter dawn

Treatment	Fruit weight (g)	Yield per plant (g)	Estimated yield per ha (q.)
Control (water)	14.84±0.31 ^d	324.42±11.49 ^e	180.23±6.38 ^e
Salicylic acid (SA)			
SA (100 ppm)	15.86±0.01 ^{cd}	358.71±0.57 ^e	199.29±0.32 ^e
SA (200 ppm)	16.07±0.42 ^{bcd}	429.02±19.40 ^d	238.34±10.78 ^d
SA (300 ppm)	16.55±0.07 ^{bcd}	441.40±3.23 ^d	245.22±1.79 ^d
Bulk chitosan			
BCH (100 ppm)	16.33±0.65 ^{bcd}	440.04±15.61 ^d	244.47±8.67 ^d
BCH (400 ppm)	16.99±0.05 ^{bc}	456.17±3.46 ^{cd}	253.43±1.92 ^{cd}
BCH (800 ppm)	17.21±0.29 ^{bc}	467.21±5.61 ^{cd}	259.56±3.12 ^{cd}
BCH (1200 ppm)	17.32±0.47 ^{bc}	479.±14.37 ^{cd}	266.12±7.98 ^{cd}
CS-SA NFs			
CS-SA NF (100 ppm)	17.10±0.13 ^{bc}	464.69±1.76 ^{cd}	258.16±0.98 ^{cd}
CS-SA NF (400 ppm)	17.78±0.58 ^{bc}	519.02±29.14 ^c	288.34±16.19 ^c
CS-SA NF (800 ppm)	20.02±0.46 ^a	659.65±0.44 ^a	366.47±0.25 ^a
CS-SA NF (1200 ppm)	17.88±0.42 ^a	595.21±0.45 ^b	330.67±0.25 ^b

The yield parameters were recorded at harvest. Each value is mean of triplicate and each replicate consisted of 5 plants. 1st foliar was done at 30 days and 2nd at 60 days after transplanting. Mean ± SE followed by same letter is not significantly different at $p = 0.05$ as determined by Tukey- Kramer HSD. BCH represents bulk chitosan dissolved in 0.5% acetic acid

Table 3. Effect of chitosan-salicylic acid NFs on quality attributes of strawberry cv. winter dawn

Treatment	TSS (°Brix)	Ascorbic acid (mg/100g)	Total sugar (%)
Control (water)	5.22±0.11 ^g	54.17±1.16	4.67±0.09 ^g
Salicylic acid (SA)			
SA (100 ppm)	6.16±0.00 ^{ef}	56.13±0.04	5.12±0.00 ^{fg}
SA (200 ppm)	6.23±0.16 ^{def}	57.06±1.52	5.26±0.14 ^{efg}
SA (300 ppm)	6.48±0.02 ^{def}	58.43±0.24	5.46±0.02 ^{def}
Bulk chitosan			
BCH (100 ppm)	6.23±0.02 ^{def}	57.27±0.24	4.66±0.02 ^g
BCH (400 ppm)	6.71±0.02 ^{cde}	59.58±0.18	6.27±0.02 ^{bc}
BCH (800 ppm)	6.87±0.11 ^{bcd}	59.62±1.02	6.38±0.10 ^{bc}
BCH (1200 ppm)	7.34±0.20 ^{abc}	60.26±1.66	6.73±0.18 ^{ab}
CS-SA NFs			
CS-SA NF (100 ppm)	5.97±0.04 ^f	56.44±0.42	5.85±0.04 ^{cde}
CS-SA NF (400 ppm)	6.44±0.20 ^{def}	56.09±1.83	6.01±0.19 ^{cd}
CS-SA NF (800 ppm)	7.54±0.17 ^a	62.91±1.45	7.30±0.16 ^a
CS-SA NF (1200 ppm)	7.43±0.17 ^{ab}	60.75±1.43	7.07±0.16 ^a

The quality parameters were recorded at harvesting. Each value is mean of triplicate and each replicate consisted of 5 samples. 1st foliar was done at 30 days and 2nd at 60 days after transplanting. Mean ± SE followed by same letter is not significantly different at $p = 0.05$ as determined by Tukey- Kramer HSD. BCH represents bulk chitosan dissolved in 0.5% acetic acid

The data regarding quality parameters revealed that maximum TSS was found in CS-SA NFs 800 ppm (7.54 °Brix) followed by CS-SA NFs 1200 ppm (7.43 °Brix) and BCH 1200 ppm (7.34 °Brix) as compared to control (5.22 °Brix) and rest of treatments. The data showed that ascorbic acid content in strawberry fruit significantly higher in CS-SA NFs 800 ppm (62.91 mg/100g) followed by CS-SA NFs 1200 ppm (60.75 mg/100g) and BCH 1200 ppm (60.26 mg/100g) as compared to control (54.17 mg/100g) and other treatments. The maximum total sugar content was also observed in CS-SA NFs 800 ppm (7.30 %) followed by CS-SA NFs 1200 ppm (7.07 %) as compared to BCH 1200 ppm (6.73 %) and lowest in control (4.67 %) (Table 3).

4. DISCUSSION

Plant height and number of branches were recorded after twice foliar spray (30 and 60 days after transplanting) of CS-SA NFs, SA, BCH and water (control). Overall, positive effect was observed specially in 800-1200 ppm concentrations of CS-SA NFs. On the basis of mean value CS-SA NFs was more effective growth promotor. Application of CS-SA NFs significantly increased plant height and number of branches as compared to control. Earlier this formulation was tested on wheat and maize [9,10]. In earlier study, salicylic acid-chitosan nanoparticles (SA-CS NFs) were developed where chitosan matrix slowly released the

encapsulated SA for its sustained availability to growing plant. Slow release of SA comprehensively stimulated plant antioxidant strength, protected maize crop from disease and improved yield as compared with SA [9]. The bio-efficacy of the developed CS-SA NF was evaluated on yield attributes in strawberry. In agreement with earlier findings, foliar application of CS-SA NFs (800-1200 ppm) significantly increased photosynthesis and vegetative growth which ultimately led to higher production and productivity in terms of fruit weight, fruit length, fruit volume, fruit diameter, number of fruits per plant, total yield plant⁻¹ and estimated yield ha⁻¹. Quality attributes viz. TSS, ascorbic acid, reducing sugar, total sugar, were significantly higher in foliar application of CS-SA NFs (800-1200 ppm). Previous studies confirms that a slow releasing SA from SA-CS NPs as compared with bulk SA considerably increases source-activity and sink- strength. Despite the prominent role of slow release of SA, intrinsic growth regulating activity of chitosan due to availability of nitrogen (N) from -NH₂ group of chitosan and phosphorus (P) from tri-sodium- polyphosphate may also contributes to plant growth, yield and ultimately the fruit quality.

5. CONCLUSION

Response of CS-SA NF in present study due to slow and sustained release of SA to maintained cellular redox, plant growth activity of chitosan and availability of N via -NH₂ group of chitosan.

We concluded that developed CS-SA NF can be tested at farmers field and other fruits crops for its efficacy.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Wange RS, Kzlogoz GA. Effect of bio-fertilizer on growth, yield and quality of strawberry. *Annals of Agricultural Science Mosthohor*. 1998;43:247-254.
2. Thakur Y, Chandel JS, Verma P. Effect of plant growth regulators on growth, yield and fruit quality of strawberry (*Fragaria x ananassa Duch.*) under protected conditions. *Journal of Applied and Natural Science*. 2017;9(3):1676-1681.
3. Anonymous. Horticulture database, government of india, ministry of agriculture and farmers welfare, horticulture statistics division. national horticulture board, Gurgaon (Haryana); 2020.
4. Darnell RL, Cantliffe DJ, Kirschbaum DS, Chandler CK, Janick J. The physiology of

- flowering in strawberry. *Horticultural Reviews*. 2010;28:325-345.
5. Hancock J, Erez A. Strawberries. *Temperate fruit crops in warm climate*; 2000.
6. Rahman M, Mukta JA, Sabir AA, Gupta DR, Mohi-Ud-Din M, Hasanuzzaman M, Miah MG, Rahman M, Islam MT. Chitosan biopolymer promotes yield and stimulates accumulation of antioxidants in strawberry fruit. *The Public Library of Science*. 2018;13(9): 0203769.
7. Aghaeifard F, Babalar M, Fallahi E, Ahmadi A. Influence of humic acid and salicylic acid on yield, fruit quality and leaf mineral elements of strawberry (*Fragaria x Ananassa Duch.*) cv. Camarosa. *Journal of Plant Nutrition*. 2016;39(13): 1821-1829.
8. Lane JM, Eynon L. Determination of reducing sugars by means of Fehling's solution with methylene blue as internal indicator. *Journal of the Society of Chemical Industry*. 1965;42: 3777.
9. Kumaraswamy RV, Kumari S, Choudhary RC, Sharma SS, Pal A, Raliya R, Biswas P, Saharan V. Salicylic acid functionalized chitosan nanoparticle: A sustainable biostimulant for plant. *International Journal of Biological Macromolecules*. 2019;123: 59-69.
10. Kadam PM, Prajapati D, Kumaraswamy RV, Kumari S, Devi KA, Pal A, Sharma SK, Saharan V. Physio-biochemical responses of wheat plant towards salicylic acid-chitosan nanoparticles. *Plant Physiology and Biochemistry*. 2021;162:699-705.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/124686>