



## **Isolation, Computational Characterization and Structure Prediction of Chitinase Gene-*Bbchit* of *Beauveria bassiana***

**N. S. Nandhan<sup>a</sup>, K. K. Kumar<sup>a\*</sup>, R. Deepa Sankari<sup>b</sup>, V. Balasubaramani<sup>a</sup> and M. Murugan<sup>c</sup>**

<sup>a</sup> Department of Plant Biotechnology, Tamil Nadu Agricultural University, Coimbatore, Tamil Nadu, India.

<sup>b</sup> Department of Plant Pathology, Tamil Nadu Agricultural University, Coimbatore, Tamil Nadu, India.

<sup>c</sup> Department of Agriculture Entomology, Tamil Nadu Agricultural University, Coimbatore, Tamil Nadu, India.

### **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/IJPSS/2022/v34i2231470

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/91266>

**Original Research Article**

**Received 17 June 2022**  
**Accepted 24 August 2022**  
**Published 27 August 2022**

### **ABSTRACT**

*Beauveria bassiana* is an entomopathogenic fungus that is used as a biopesticide for the control of many insect pests in agriculturally important crops. *B. bassiana* produces many extracellular hydrolytic enzymes that degrade the insect cuticle. Among them, the fungal exochitinase mainly assists in the infection by degrading the chitin present in the insect cuticle. To characterize the exochitinase from *B. bassiana*, the chitinase gene (*Bbchit*) of 1050bp was amplified from the genomic DNA through PCR using gene-specific primers. This chitinase gene was cloned into the pJET1.2 cloning vector, confirmed through sequencing. The sequence (GI; OP114061) has been deposited in the NCBI database and used for bioinformatics analysis. The various Physico-chemical properties of chitinase protein such as pI, EC, AI, GRAVY and instability index were predicted using primary structure analysis. Using MEGA11 software, the translated chitinase protein- *Bbchit* showed that it closely resembles with the QJE37897.1 (chitinase Chit37). Primary and secondary structure analysis helps to characterize the functionality of chitinase protein (*Bbchit*). The 3D structure of *B. bassiana* was predicted using Robetta, an abinitio prediction method and

\*Corresponding author;  
E-mail: nandhansriram98@gmail.com;

confirmed through in-silico using the SAVESv6.0 (PROCHECK) server. The isolated exochitinase gene from *B. bassiana* has been characterized for use as a bio-control agent for the control of insect pests in crop plants.

**Keywords:** *Bbchit*; *Beauveria bassiana* (*Bb*); chitinase; physico-chemical; protein structure prediction.

## 1. INTRODUCTION

*Beauveria bassiana* (Balsamo) Vuillemin belongs to the class Sordariomycetes and is one of the well-known genera of entomopathogenic fungi, used as a biopesticide to control a wide range of plant-insect pests [1]. Agostino Bassi di Lodi (1835) discovered that *B. bassiana* was responsible for the white muscardine disease of silkworms (*Bombyx mori*). As cosmopolitan fungi found on infected insects in both temperate and tropical regions, *B. bassiana* habitats range from desert soils to forest and cultivated soils [2]. *B. bassiana* secretes many extracellular enzymes including chitinase, proteases, lipases, amylase, laccase and others [3]. Chitin, a homopolymer chain of N-Acetylglucosamine (GlcNAc) linked by  $\beta$ -1,4glucosidic linkage, is cleaved by chitinase, an enzyme that has been extensively studied as a pest control agent [4]. In addition, *B. bassiana* also produces a plethora of biologically active secondary metabolites such as polyketides, peptides and oxalic acid, all of which are useful in parasitizing and killing hosts [5].

Earlier studies reported the isolation of an endochitinase gene (*Bbchit1*) from *B. bassiana* [6]. Increased aphid pathogenicity was observed when the *Bbchit1* gene was overexpressed in transgenic *B. bassiana* [7]. The transformation of the *B. bassiana* chitinase gene into Chinese white poplar (*Populus tomentosa*) resulted in resistance to infection by the fungal pathogen, *Cytospora chrysoperma* [8]. The genome of *B. bassiana* strain ARSF2860 was sequenced for a better understanding of pathogenesis and interactions between insects and plants [9] and high throughput RNA-seq transcriptomic analysis revealed that *B. bassiana* can survive in a variety of environmental niches by stimulating well-defined gene sets. One integrated crop pest management tool employs *B. bassiana* as a mycoinsecticide, which aids research into improving fermentation and formulation technologies [10].

In our present investigation, we have taken up to isolate and characterize an insect cuticle degrading chitinase gene, *Bbchit* from the entomopathogenic fungus *B. bassiana*. Using the

various bioinformatics tools, the Physico-chemical properties, as well as the primary, secondary and tertiary structures of *B. bassiana* chitinase protein was determined. Phylogenetic analysis was done to study the evolutionary comparisons with other accessions of *B. bassiana* chitinase. This computational study will aid in understanding the biological function of the chitinase protein and the mechanism of chitin degradation.

## 2. MATERIALS AND METHODS

### 2.1 Isolation of Genomic DNA from *Beauveria bassiana*

The fungal genomic DNA from *B. bassiana* was isolated by the CTAB method [11]. One gram of the *B. bassiana* mycelial mat was homogenized to a powder with a pre-chilled pestle and mortar using liquid nitrogen (-196°C). About 700 $\mu$ l of pre-warmed CTAB buffer containing  $\beta$ -mercaptoethanol was added to the powdered sample. Then the mixture was transferred to a 2 ml microfuge tube and incubated at 65°C for 40 min in a water bath (mixed by inversion at 10 min intervals). It was then centrifuged at 4°C for 10 min at 13000 rpm. The supernatant was transferred to a new microfuge tube and an equal volume of phenol: chloroform: isoamyl alcohol (25:24:1) mixture was added. After thorough mixing, the mixture was centrifuged at 4°C for 10 min at 13000 rpm and the aqueous layer was transferred to a new microfuge tube. To the above mixture, an equal volume of ice-cold isopropanol was added and incubated at -20°C overnight. The DNA was pelleted by centrifuging at 12000 rpm for 10 min at 4°C. After discarding the supernatant, the pellet was washed twice with ice-cold 70% ethanol followed by centrifuging at 12000 rpm for 5 min at 4°C. Nuclease-free water (50  $\mu$ l) was used to resuspend air-dried DNA pellets.

### 2.2 Amplification of *B. bassiana* Chitinase Gene

The chitinase-specific primers *Bbchit\_F* (5'-ATGGCTCCTTTTCTTCAAACCCAGCC-3') and *Bbchit\_R* (5'-

TTACGCAGTCCCCAAAGTCCCCT-3') were designed using the *B. bassiana* nucleotide sequence (NCBI Accession No. XM008601414.1). The PCR reactions contained 50 ng of *B. bassiana* genomic DNA, 1 µmol forward and reverse primers, 1X PCR buffer, 1 unit of Taq DNA polymerase (Takara), 0.25 mM dNTPs and nuclease-free water to make up the volume of 40 µl. The amplification was performed by following PCR profile; initial denaturation (5 min at 95°C), followed by 30 cycles (denaturation (45 sec at 95°C), annealing (45 sec at 55°C), extension (45 sec at 72°C)) and final extension (10 min at 72°C). The PCR products were then resolved in 1.0% agarose gel and the amplified product of ~1.0 kbp was eluted, purified and used for cloning.

### 2.3 Cloning of *Bbchit* Gene of *B. bassiana* into pJET1.2 Vector

The purified PCR product was cloned into pJET1.2/blunt cloning vector. The ligation reaction contains 50 ng of purified PCR product, 2X reaction buffer, 1 µl of pJET1.2 cloning vector (50 ng), 1 unit of T<sub>4</sub> DNA ligase and nuclease-free water to make up the volume of 20 µl. The cocktail is vortexed and centrifuged for 3-5 seconds and incubated at 22 °C for 20 min. The entire ligated product is used for transformation into *E.coli*- DH5α competent cells.

### 2.4 Sequence Retrieval and Phylogenetic Analysis

For computational analysis, the amino acid sequences of eight chitinases for different accessions of *B. bassiana* with 100% query coverage were obtained in FASTA format from the NCBI database [12]. MEGA11 software was used to generate the phylogenetic tree for the retrieved sequences and their evolutionary studies were determined using the UPGMA (Unweighted Pair Group Method with Arithmetic Mean) method.

### 2.5 Physico-Chemical Characterization of Chitinase

The Protparam [13] tool was used to calculate the Physico-chemical parameters of deduced chitinase protein sequence. Isoelectric point (pI), the total number of amino acids (+ve and -ve charged residues), instability index (II), molecular weight, extinction coefficient (EC), aliphatic index (AI) and grand average hydropathy (GRAVY) were all calculated [14].

### 2.6 Functional and Secondary Structure Analysis

The Prosite, Pfam, and SOSUI servers were used to perform functional annotation of the chitinase protein. The length of amino acid residues of proteins with specific profiles and patterns was recorded in ExPASy-prosite [15,16], a database of protein families and domains. The conserved domain of *B. bassiana* chitinase was checked using Pfam [17].

Self-Optimized Prediction Method with Alignment (SOPMA) server [18,19] was used to predict the secondary structure of chitinase with default parameters such as Similarity threshold 8 and Window width 17. It computes the content of α-helix, β-sheets, turns, random coils, and extended strands based on the amino acid sequence.

### 2.7 Molecular Modelling of Chitinase

The protein three-dimensional structure of *B. bassiana* full-length chitinase was not available in the PDB. As a result, the tertiary structure of the protein- chitinase was modeled using the Robetta web server [20]. The quality of the predicted 3D model was validated by the SAVES v6.0 server to study the overall stereo-chemical property of protein i.e. to evaluate the energetically allowed regions.


## 3. RESULTS AND DISCUSSION

### 3.1 PCR Amplification of the *Bbchit* gene of *Beauveria bassiana*

The *Bbchit* gene was amplified by using designed chitinase gene-specific primers. The amplified fragment of the *Bbchit* gene when analyzed by agarose gel electrophoresis (Fig. 1), the expected size *Bbchit* gene was observed. The amplified DNA was eluted and cloned into the pJET1.2 vector (Thermo Scientific).

### 3.2 Sequence Analysis of the *Bbchit* Gene

To verify the nucleotide sequence of a cloned PCR product in the pJET1.2 vector. The plasmid DNA was isolated and then sequenced with both vector-specific and gene-specific primers. The sequence of the cloned *Bbchit* gene is 1050 bases (Annexure 1).




**Fig. 1. PCR amplified *Bbchit* gene from *B. bassiana***

The cloned exochitinase gene sequence was deposited in the NCBI database (accession number: OP11406). This gene sequence was used for further bioinformatics analysis. The translated protein sequence revealed an open reading frame encoding the exochitinase gene of 1047 bases. Performing an NCBI protein blast search with translated protein revealed that one of the *B. bassiana* gene deposited from Colombia had 100% sequence homology. However, at the nucleotide level, the isolated chitinase *Bbchit* gene showed 98.38 % sequence similarity with accession no. MN871410.1. The dissimilarity in nucleotide sequence helped to proceed to check the evolutionary relationship at the protein level.

### 3.3 Sequence Retrieval and Phylogenetic Analysis

Using the Expsasy tool – translate [21], the nucleotide sequence of *B. bassiana* chitinase (GI; OP114061) was translated to amino acid

sequence. The translated sequence was tested for similarity using BLASTP. The full-length translated protein sequences of *B. Bassiana* were used for the phylogenetic analysis with other *B. bassiana* chitinase genes deposited in NCBI database and they are QJE37897.1 (chitinase Chit37), ADP44708.1 (chitinase), ACF32998.1 (chitinase), XP 008599636.1 (chitinase-like protein), AEG21075.1 (chitinase), ACZ28129.1 (endochitinase), UQK86746.1 (chitinase) and AAN41259.1 (chitinase). The translated chitinase protein- *Bbchit* closely resembles the QJE37897.1 (chitinase Chit37) and was distantly related (out cluster) to the ACZ28129.1 (endochitinase).Based on the amino acid sequences of the chitinase protein from different *B. bassiana* strains, the clusters with the highest similarity were observed in comparison analysis (Fig. 2). Except the ACZ28129.1 (endochitinase) strain, translated chitinase and other chitinase strains belong to the exochitinase subgroup.



**Fig. 2. Phylogenetic analysis of chitinase from different accessions of *B. bassiana*: In this Fig., a and b are 2 clusters, c is the out-group, p and q are closely related**

### 3.4 Physico-Chemical Characterization of Chitinase

Physico-chemical parameters analysis done with translated *B. bassiana* chitinase enzyme has a molecular weight of 36.78kDa. The total number of amino acids is 348 residues, of which 26 are negatively charged and 23 are positively charged. The chitinase enzyme has a calculated pI value of 5.94, which have acidic proteins and plays an important role in chitinase activity. The instability index value is 30.14, indicating that the protein is stable (index value is less than 40) [22] and has an *in vivo* half-life of more than 20 hours in yeast, >10 hours in *Escherichia coli* and 30 hours in mammalian reticulocytes. The aliphatic index is the relative volume occupied by the aliphatic side chains (alanine, valine, isoleucine and leucine) and it predicts protein stability over a wide temperature range. Chitinase protein has an aliphatic index of 84.17 and is predicted to be thermostable based on the report of Ikai, 1980 [23]. The GRAVY value of -0.027 indicates that the chitinase protein is hydrophilic [14].

### 3.5 Functional and Secondary Structure Analysis


The pattern and profile hit of the proteins were analyzed using ExPASy Prosite. The chitinase protein has 18 active sites, including the N-myristoylation site (14 regions), Casein kinase II phosphorylation site (2 regions), Protein kinase C phosphorylation site (single) and N-glycosylation site (single). The domain identified by profile for the chitinase enzyme was the GH 18 catalytic domain (position; 41-348 residue), with glutamic acid as the active site at residue 163. The chitinase enzyme was classified as a glycosyl

hydrolase using the Pfam tool (GH 18). The glycosyl hydrolase-18 family's chitinase is a hydrolytic enzyme that cleaves the  $\alpha$ -1,4-bond, releasing oligomeric, dimeric (chitobiose), or monomeric (N-acetyl glucosamine, GlcNAc) products. The SOSUI online server indicates that the chitinase protein is water soluble.

The SOPMA tool is used to find the distribution of amino acids as helix, strand, turn, or coil. The results for the chitinase protein show that random coils dominate secondary structure elements, followed by  $\alpha$ -helix, extended strand and  $\beta$ -turns for all sequences (Table 1). High random coil studies lead to intrinsic confirmations that correlate with low energy confirmations and have implications for protein structure prediction and design.

### 3.6 3-D Modeling and Validation


The tertiary structure of proteins will aid in the study of the protein's function and active sites. Robetta online server was used to create a 3D model of the chitinase protein. Using the PyMOL molecular visualization tool, the predicted model for chitinase was visualized (Fig. 3). The SAVES server was used to evaluate the predicted model. The stereo-chemical quality of the model was classified using the PROCHECK tool. The Ramachandran plot (Fig. 4) depicts the statistical distribution of the possible combinations of backbone dihedral angles  $\phi$  and  $\psi$  for all the amino acid [24]. The total number of residues distributed in the most distributed region is approximately 91.5 %, indicating that the modeled structure is reliable and of high quality (Table 2).



**Fig. 3. 3D structure of chitinase protein visualized through PyMOL: The cyan color represents the  $\alpha$ -helix, the red color represents the  $\beta$ -sheets and the magenta color represents random coils**

**Table 1. Secondary structure elements are calculated by SOPMA**

Structures	$\alpha$ - helix	Extended strand	$\beta$ - turn	Random coil
No. of residues	108	69	22	149
Distribution	31.03 (%)	19.83 (%)	6.32 (%)	42.82 (%)



**Fig. 4. Ramachandran plot showing the phi -psi torsion angles for all the residues of the chitinase protein: the distribution of amino acid residues to  $\beta$ -sheets (+psi, -phi), right-handed  $\alpha$ -helix (-psi, -phi) and left-handed  $\alpha$ -helix (+phi, +psi)**

**Table 2. Ramachandran plot calculation with Procheck program**

Ramachandran plot statistics	Distribution
Residues in most favoured regions	91.5 (%)
Residues in additionally allowed regions	7.1 (%)
Residues in generously allowed regions	0.0 (%)
Residues in disallowed regions	1.4 (%)

#### 4. CONCLUSION

Exochitinase gene of *B. bassiana* was amplified, cloned and sequenced. The translated chitinase enzyme shows that it was having 100% sequence homology with the QJE37897.1 (chitinase Chit37). The primary structure of chitinase protein reveals that it is acidic. According to the functional analysis, the protein is stable, hydrophilic, and soluble in water. The Ramachandran plot validated the 3D model of the chitinase protein created by Robetta, which has a 91.5% distribution of amino acid residues in the most favoured region. This research on *B. bassiana* chitinase would be useful for *in vitro* molecular studies such as to know how the chitinase enzyme help to degrade the chitin

polymer of many insects and fungal strains to act as a bio-control agent.

#### ACKNOWLEDGEMENTS

The authors express sincere thanks to the Department of Plant Biotechnology, Tamil Nadu Agricultural University, Coimbatore for providing facilities to conduct the research. The authors express their sincere gratitude to ICAR- ITCC for providing *Beauveria bassiana* cultures for the study.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Akello J, Dubois T, Coyne D, Kyamanywa S. The effects of *Beauveria bassiana* dose and exposure duration on colonization and growth of tissue cultured banana (*Musa* sp.) plants. *Biological Control*. 2009; 49(1):6-10.
2. Keswani C, Singh SP, Singh HB. *Beauveria bassiana*: Status, mode of action, applications and safety issues. *Biotech Today*. 2013;3(1):16-20.
3. Amobonye A, Bhagwat P, Pandey A, Singh S, Pillai S. Biotechnological potential of *Beauveria bassiana* as a source of novel biocatalysts and metabolites. *Critical Reviews in Biotechnology*. 2020;40(7): 1019-1034.
4. Songjang K, Donchai T, Chaiyawat P, Meyer C. Cloning and expression of chitinase gene isolated from insect pathogenic fungi, *Beauveria bassiana* in *Escherichia coli*. *Chiang Mai J Sci*. 2006;33(3):347-355.
5. Wang H, Peng H, Cheng, P, Gong, M. The toxins of *Beauveria bassiana* and the strategies to improve their virulence to insects. *Frontiers in Microbiology*. 2021; 2375.
6. Fang W, Leng B, Xiao Y, Jin K, Ma J, Fan Y et al. Cloning of *Beauveria bassiana* chitinase gene Bbchit1 and its application to improve fungal strain virulence. *Applied and Environmental Microbiology*. 2005; 71(1):363-370.
7. Rajasekhar P, Kalidas P, Rao KRSS. Cloning and expression of Bbchit1 gene of *Beauveria bassiana*. *Open Entomology Journal*. 2010;4:30-35.
8. Jia Z, Sun Y, Yuan L, Tian Q, Luo K. The chitinase gene (Bbchit1) from *Beauveria bassiana* enhances resistance to *Cytospora chrysosperma* in *Populus tomentosa* Carr. *Biotechnology letters*. 2010;32(9):1325-1332.
9. Xiao G, Ying SH, Zheng P, Wang ZL, Zhang S, Xie XQ, et al. Genomic perspectives on the evolution of fungal entomopathogenicity in *Beauveria bassiana*. *Scientific Reports*. 2012;2(1):1-10.
10. Mascarin GM, Jaronski ST. The production and uses of *Beauveria bassiana* as a microbial insecticide. *World Journal of Microbiology and Biotechnology*. 2016; 32(11):1-26.
11. Carter-House D, Stajich JE, Unruh S, Kurbessoian T. Fungal CTAB DNA Extraction. *Protocols*. Io; 2020.
12. Available:<https://www.ncbi.nlm.nih.gov/prot/ein/>
13. Available:<https://web.expasy.org/protparam/>
14. Kyte J, Doolittle RF. A simple method for displaying the hydropathic character of a protein. *Journal of Molecular Biology*. 1982;157(1):105-132.
15. Available:<https://prosite.expasy.org/>
16. Falquet L, Pagni M, Bucher P, Hulo N, Sigrist CJ, Hofmann K, et al. The PROSITE database, its status in 2002. *Nucleic Acids Research* 2002;30(1):235-238.
17. Available:<https://pfam.xfam.org/>
18. Available:[https://npsaprabi.ibcp.fr/cgi-bin/npsa\\_automat.pl?page=/NPSA/npsa\\_sopma.html](https://npsaprabi.ibcp.fr/cgi-bin/npsa_automat.pl?page=/NPSA/npsa_sopma.html)
19. Ashokan KV, Mundaganur DS, Mundaganur YD. Catalase: Phylogenetic characterization to explore protein cluster. *J of Res in Bioinfo*. 2010;1:1-8.
20. Available:<https://rosetta.bakerlab.org/queue.php>
21. Available:<https://web.expasy.org/translate/>
22. Gasteiger E, Hoogland C, Gattiker A, Wilkins MR, Appel RD, Bairoch A. Protein identification and analysis tools on the ExPASy server. *The Proteomics Protocols Handbook*. 2005;571-607.
23. Ikai A. Thermostability and aliphatic index of globular proteins. *The Journal of Biochemistry*. 1980;88(6):1895-1898.
24. Spencer RK, Butterfoss GL, Edison JR, Eastwood JR, Whitelam S, Kirshenbaum, et al. Stereochemistry of polypeptoid chain configurations. *Biopolymers*. 2019;110(6): e23266.



### Annexure 1. Sequence of the *Bbchit* gene (GI: OP11406)

TGGCTCCTTTTCTTCAAACCAGCCTCGCGCTCCTTCCATTGTTGGGCTTTCCAACCATGGTCAGC  
GCCTCGCCATTGGCGCCGCGAGCCGACACCTGCGCAACCAAGGGCCGGCCGGCCGCAAAGT  
GCTCCAGGGCTACTGGGAGAAGTGGGACGGTGCCAAGAACGGCGTGCACCCTCCGTTTGGCTG  
GACGCCATCCAAAACCCCGACATTGCAAGCACGGCTACAACGTCATCAATGCTGCCTTTCCC  
ATCATCCAGCCCGACGGCACC GCGCTCTGGGAGGACGGCATGGACACGGGCGTCAAGGTGGC  
GAGCCCGGCCGACATGTGCGAGGCCAAGGCAGCGGGCGCCACCATCTTGATGTCGATTGGCG  
GCGCTACTGCGGCCATTGACCTGAGCTCGTCGGCTGTGGCTGACAAGTTTGTCTCGACCATTGT  
GCCGATTCTGAAAAAGTACAACCTTTGACGGCATTGATATCGACATTGAATCCGGCCTCACAGGCA  
GCGGAAACATAAACACCCTGTCCACCTCGCAGACCAACCTGATTAGAATCATTGACGGCGTTCTC  
GCGCAGATGCCCGCCAACCTTTGGCTTGACCATGGCGCCAGAGACTGCCTACGTTACCGGTGGG  
ACGATTACGTACGGATCAATCTGGGGCTCTTACCTCCCCATCATCAAAAAGTACCTGGACAATGG  
TCGTCTCTGGTGGCTCAACATGCAGTACTACAATGGCGAAATGTACGGCTGCTCCGGCGACTCG  
CACAAGGCCGGTACTGTGCAAGGGTTTCGTTGCTCAGACCGACTGCCTGAACAAGGGACTTAGTA  
TTCAGGGCGTGACAATCACGATTCCCTATGACAAGCAAGTGCCTGGCCTTCCTGCCAGCCTGG  
GGCTGGCGGGCGCCACATGTCCCCGTCCAACGTGGCGCAAGTTCTCTCCCACTACAAGGGCGC  
TTTGAAGGGATTGATGACTTGGTCTCTGAACTGGGACGGCTCCAAGAATTGGACATTTGGCGAC  
AATGTCAAGGGGACTTTGGGGACTGCGTAA

© 2022 Nandhan et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/91266>