



# Impact of Early Rehabilitation in a Complex Case of Non-union of Tibial Plafond Fracture with Osteosynthesis Associated Infection – A Case Report

Harsh Nathani <sup>a#</sup>, Medhavi V. Joshi <sup>at</sup> and Pratik A. Phansopkar <sup>at\*</sup>

<sup>a</sup>Department of Musculoskeletal Physiotherapy, Ravi Nair Physiotherapy College, Datta Meghe Institute of Medical Sciences, Sawangi Meghe, Wardha, Maharashtra, India.

## Authors' contributions

All the authors made the best contribution for the concept, assessment and evaluation, data acquisition, analysis and interpretation of data.

## Article Information

DOI: 10.9734/JPRI/2021/v33i51B33511

### Editor(s):

(1) Dr. Vasudevan Mani, Qassim University, Saudi Arabia.

### Reviewers:

(1) Alex Guedes, Universidade Federal da Bahia, Brazil.

(2) Isiaka Adio Hassan, Lagos State Polytechnic, Nigeria.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:

<https://www.sdiarticle5.com/review-history/76727>

Case Report

Received 09 September 2021

Accepted 19 November 2021

Published 24 November 2021

## ABSTRACT

**Background:** Fractures of distal end of tibia associated with soft tissue injuries and fracture of distal fibular end are very complex and forms a total of 1-2% of all fracture of lower limb. These fractures are widely termed as plafond fractures.

**Case Presentation:** A 26- year-old male, a follow up case, gave a history of road traffic accident following which he underwent corticotomy and application of external Ilizarovring fixator. At present due to non-union of the fracture segments patient got readmitted after a year. Further management through a three-step surgical approach was carried out. Rehabilitation program began from post-operative day 1 and was continued for a period of three weeks.

**Investigations:** On the day of examination, the patient's pain was severe on movement with presence of disuse trophy of lower limb musculature of the affected extremity. Ranges on the right

<sup>#</sup> BPT Student

<sup>†</sup> Resident

<sup>‡</sup> Associate Professor & HOD

<sup>\*</sup> Corresponding author: E-mail: drpratik77@gmail.com;

lower limb at all joints were reduced due to pain. The X-ray showed presence of 9-hole recon plate fixed distally over talus and proximally to tibia.

**Management:** Physiotherapeutic intervention began with educating the patient and the caregivers about the condition, the precautions to be taken, the expected time of healing and extent of healing. The exercise program was based on the principles of variability and individuality. The protocol was changed weekly with the observed progression in the patients range, muscles strength and ability to perform more challenging in bed activities.

**Conclusion:** Early rehabilitation in complex cases of tibial plafond fracture facilitates the process of healing as well as maintain the patients level of functioning by maintain muscle properties. Post-operative complications are also reduced.

*Keywords: Tibialfracture; osteosynthesis; rehabilitation.*

## 1. INTRODUCTION

Tibia and fibula are the long bones of the lower limb, forming a tibiofibular syndesmosis. Fracture of these long bones due to high energy axial compression is also known as tibia plafond fractures and can have an associated soft tissue involvement and extension of fracture into fibula [1]. These fractures are rare and account for only 1-10 % of total fractures of lower limbs. The type 3 fracture of this is the most severe with multiple fracture fragments and metaphysical impaction [2].

These fractures are managed surgical in 4 phases, beginning from restoration of fibular length, followed by autologous bone graft for filling the defect and ending by insertion of buttress plate on distal aspect of tibia [3,4]. Even with advancement in the surgical approaches used long term complication of non- union still persists and therefore the fracture was also termed as not "amenable" to surgery [2].

Osteosynthesis associated infection (OAI) is one of the most common complication post- surgical management of these fractures indicated by infection in the presence of an implant. OAI is difficult to treat due to presence of instable fracture site, open wound or trauma and need for preservation of cartilage [5].

## 2. CLINICAL PRESENTATION

A 26-year-old male, a follow up operated case came to the hospital on 1/10/21 for grade III B fracture of distal tibiofibular joint of right leg. The patient presents with the history of road traffic accident 2 years back on 16/09/19 following which he sustained injuries to right leg. There was an open wound with excessive blood loss associated with pain was sudden in onset. The pain was progressive and continuous in nature.

There was no history of head injury, loss of consciousness or chest trauma. Patient was then by the bystanders brought to the hospital where he was managed with external fixator application on the same day. The patient was then discharged and since then, at home the patient had a non- weight bearing ambulatory functional status. The patient was then re-admitted on 1/10/19 for further wound management. The x ray revealed non- union of the distal tibia fibula fracture segment for which he was operated on 4/10/21 where external ilizarov ring fixator was removed and open reduction internal fixation with plate osteosynthesis. The second operation was conducted on the same day where bone graft from right iliac fossa and skin graft from left thigh was taken. Since then the patient is referred for physiotherapy for further rehabilitation.

## 3. CLINICAL FINDINGS

### 3.1 On Observation

Patient was lying in supine lying with hands by the side, right lower extremity was in elevation with pillows underneath and in slight external rotation and in 5 degrees of plantarflexion. Wound dressing was present from the balls of the feet upto mid knee joint line. Quadriceps muscle wasting could be seen indicating disuse atrophy.

### 3.2 On Palpation

Distal pulses and tenderness at the ankle could not be assessed due to dressing. Tenderness of grade 2 at the right iliac region at the suture site was present indicated by pain and flinching by the patient on palpation. Swelling over the exposed parts of left lower extremity was absent. Muscle wasting of quadriceps was quantitatively confirmed by a difference of 3.5 cm during limb girth measurement.

### 3.3 On Examination

Pain: On numeral pain rating scale during movement 6/10 (passive dorsiflexion and assisted hip flexion) and at rest 2/10.

Range of motion: All the ranges for right lower limb at hip knee and ankle were with normal functional ranges. There was no active dorsiflexion present and rest of the ranges at knee and hip were restricted due to pain giving an empty end feel.

Manual Muscle Strengthening: Manual muscle testing for left side for all the major muscle groups was 3+ ( Fair ) indicating ability to perform complete range of motion against gravity with mild resistance.

### 3.4 Investigations

Pre and post -operative X- rays and clinical images of the wound are as follow:

**Table 1. Range of motion of right affected extremity**

Joint movement on Right side	Active ( in Degrees)	Passive ( in Degrees)	End feel
Ankle Plantarflexion	5-10	5-12	Empty
Ankle dorsiflexion	0	0	-
Knee flexion	0-60	0-65	Empty
Knee Extension	60-0	60-0	Firm
Hip flexion	0	0-40	Empty
Hip abduction	0-10	0-15	Empty
Hip Adduction	Cannot be assessed		-

**Table 2. Grades of Manual Muscles Strength (MMT)**

Muscle Tested	Grade of MMT
Ankle Plantar flexors	Weak painless ( On Resisted isometric contraction
Ankle Dorsiflexors	0 ( no visible palpable contraction )
Knee extensors	Strong painless ( on resisted isometric contraction )
Hip Flexors	Could not be assessed.
Hip Abductors	2- ( partial range of motion in gravity eliminated )



**Fig. 1. Pre -Operative x-ray of right lower extremity, Lateral view. Bones exposed are tibia fibula, talus, calcaneus, navicular and metatarsals. (Orange) Arrow points at fracture segments and (red) arrow points at presence of swelling**



**Fig. 2.** Post- operative X-ray indicating the presence of nine hole recon plate (arrows) and 4.9 mm locking screw and 4.5 mm cortical screws



**Fig. 3.** X-ray is an Antero-Posterior view of Pelvis. Bones exposed are Pelvis, Sacrum and head of femur bilaterally. Arrow indicates the site of bone graft from the iliac fossa



**Fig. 4. Clinical Image of the wound. Arrow (orange) indicates the Flap taken for covering the wound taken from the proximal anterior-lateral aspect of the left below knee. Arrow (red) indicates the area where skin grafting is done**

**Table 3. Further exercise program tailor made according to the patient’s progress was as follows**

<b>Exercise type</b>	<b>Intensity Day 1-3</b>	<b>Intensity Day 4-7</b>	<b>Intensity Week 2-3</b>	<b>Rationale</b>
Ankle toe movements Bilaterally	10 repetitions X 1 set 50-70 / per day	10 repetitions x 2 set 70-100/day	10 repetitions X 2 sets 150 reps /day	To prevent edema and maintain peripheral circulation
Passive dorsiflexion Left ankle	3 repetitions X 1 set Thrice a day	3 repetitions X 1 set Every two hour	3 repetitions X 1 set Every one hour	To prevent deformity
Static Hamstring Contraction	10 repetitions 50 / per day Without hold	10 repetitions 70-80/per day With 5 seconds hold	10 repetitions 100-150 /per day with 10 second hold	To maintain muscle properties
Static Quadriceps contraction				
Static gluteal contraction				
Active Knee Flexion	Actively in pain free range	Active assisted in pain free range	Active assisted pushing slightly beyond available range	
Active assisted Hip Abduction	10 reps thrice a day	10 reps thrice a day with minimal assistance	10 reps 2 sets thrice a day	
Active assisted Straight Leg Raise	10 reps thrice a day	10 reps with 5 second hold	10 reps thrice a day with 10 second hold	
Unilateral Bridging	-	10 repetitions Without hold	10 repetitions with 5 second hold 50 reps throughout the day	

### 3.5 Management

Rehabilitation started with patient education where patient was educated about his current condition, expected time period of recovery and the precautions of wound care to avoid re-infection and bed sores. The care givers were also educated to increase the compliance towards exercise and facilitate the process of rehabilitation.

Prevention of secondary complications like, embolism and oedema was done by advising the patient to continuously perform ankle toe movements (100-150 repetition / day) and keep the limb elevated.

For the unaffected left lower extremity and bilateral upper limbs mobility and strengthening exercises were initiated from day 1 of rehabilitation. Deep breathing and pursed lip breathing exercises were taught and advised to perform every 2 hourly.

### 4. DISCUSSION

Fracture of tibiapilon are uncommon and difficult to manage and the severity depends upon position and rotation of the foot during the time of impact [6]. Open reduction internal fixation is the most common method of management with the aim of providing stability to the ankle joint for early mobility but due to more long term complication like non-union and infection, external ring fixator is the next choice of treatment [3]. A study by Maxim et al. [7] concluded that the reduction quality analyzed through 3D imaging intraoperatively was a major factor for determination of long term outcome of the surgery and rehabilitation and functional outcome. Another study suggests that most of the patient were treated through two stage, where first stage involved external fixator application for a period of few weeks to months depending on severity followed by open reduction internal fixation. It is seen that the extent of involvement of soft tissues and comminution of fracture segments is the basis for selecting the operative approach and procedure [8].

A study conducted by Faizal et al. [9] concluded that early rehabilitation post pilon fracture including isometrics, strengthening and initiating mobility within available ranges results in significant improvement statistically and a raised level of Mazur score, an outcome measure post

pilon fractures. American College Of Sports Medicine also supports the findings of the study that the absence of any forces like weight bearing as a result of long term immobilization leads to insufficiency in musculoskeletal as well as circulatory system that further delays the process of healing and worsens the prognosis [10].

The findings of this study is in line with the outcomes of research conducted by Puthoff et al. [11] that week isometric strength of lower extremity musculature, if not initiated early during rehabilitation makes it difficult for the patient to perform mobility or pat activities further in the phase of rehabilitation when initiation of standing from sitting and activation of muscles for balancing will be required.

### 5. CONCLUSION

The three-week rehabilitation program, we conclude that early initiation of range of motion, isometrics and strengthening exercises lead to a reduction in symptoms of post-operative stiffness and facilitates healing. In casewhich presents with a prolonged history of reduced mobility due to long term application of Ilizarov ring fixator where atrophy is present, such tailor-made protocols help in early recovery of the affected area and increases patients awareness towards re-infection and motivation. Variability with respect to intensity duration and type of exercise performed also plays a major role in keeping the patient motivated and compliant.

### CONSENT

Patient was informed about the study and informed consent was taken from the patient.

### ETHICAL APPROVAL

It is not applicable.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

1. Germann CA, Perron AD, Sweeney TW, Miller MD, Brady WJ. Orthopedic pitfalls in the ED: tibial plafond fractures. *The American Journal of Emergency Medicine.* 2005 May 1;23(3):357–62.

2. Luo TD, Pilson H. Pilon Fracture. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2021 [cited 2021 Oct 21]. Available:<http://www.ncbi.nlm.nih.gov/books/NBK482176/>
3. Marsh JL. Distal Tibial and Plafond Fractures. In: De Bastiani G, Apley AG, Goldberg A, editors. Orthofix external fixation in trauma and orthopaedics [Internet]. London: Springer; 2000 [cited 2021 Oct 21]:286–98. Available:[https://doi.org/10.1007/978-1-4471-0691-3\\_27](https://doi.org/10.1007/978-1-4471-0691-3_27)
4. Javdan M, Tahririan MA, Nouri M. The role of fibular fixation in the treatment of combined distal tibia and fibula fracture: A randomized, control trial. *Adv Biomed Res.* 2017 Apr 25;6:48.
5. Fang C, Wong T-M, Lau T-W, To KK, Wong SS, Leung F. Infection after fracture osteosynthesis – Part I: Pathogenesis, diagnosis and classification. *J Orthop Surg (Hong Kong).* 2017 Jan 1;25(1):2309499017692712.
6. Bone LB. Fractures of the tibial plafond. The pilon fracture. *Orthop Clin North Am.* 1987 Jan;18(1):95–104.
7. Privalov M, Euler F, Keil H, Swartman B, Beisemann N, Franke J, et al. Influence of reduction quality on functional outcome and quality of life in treatment of tibial plafond fractures: a retrospective cohort study. *BMC Musculoskeletal Disorders.* 2019 Nov 13;20(1):534.
8. Barei D, Nork S. Fractures of the tibial plafond. *Foot and Ankle Clinics of North America.* 2008;13:571-591. [Internet]. [cited 2021 Oct 21]. Available:<http://www.sciencedirect.com/science/article/pii/S1067251708000245>
9. Adam FF, Desouky AA, Abd-Elmegeed ES. Pilon fracture rehabilitation exercises program outcome. *American Journal of Nursing Research.* 2018 Jun 13;6(5):222–8.
10. ACSMs Guidelines for Exercise Testing and Prescription [Internet]. [cited 2021 Oct 21]. Available:<https://www.acsm.org/read-research/books/acsms-guidelines-for-exercise-testing-and-prescription>
11. Puthoff ML, Nielsen DH. Relationships among impairments in lower-extremity strength and power, functional limitations, and disability in older adults. *Phys Ther.* 2007 Oct;87(10):1334–47.

© 2021 Nathani et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/76727>