



Haematological Alterations in Heat-Stressed Male Wistar Rats

Onyinye Ukamaka Umeh^a
and Bruno Chukwuemeka Chinko^{a*}

^a Department of Human Physiology, Faculty of Basic Medical Sciences, College of Health Sciences, University of Port Harcourt, Port Harcourt, Nigeria.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/97945>

Original Research Article

Received: 21/01/2023

Accepted: 26/03/2023

Published: 29/03/2023

ABSTRACT

Background: Heat stress (HS) occurs due to global rising temperatures and the exposure of certain industrial workers to hot ambient temperatures. Physiological adaptability to heat stress involves long-term hemorheological modifications.

Objectives: The present study evaluated the effect of heat stress on haematological profile using heat-stressed Wistar rat models.

Materials and Methods: The study involved using twenty (20) male apparently healthy Wistar rats, with a weight range of 200-250g and an age range of 12-16 weeks. Before the study, the rats were acclimated for two weeks under standard animal husbandry conditions. To simulate heat stress (HS), a heating chamber was utilized and maintained at a temperature of $38\pm 1^\circ\text{C}$. The animals were randomly grouped into five (5), comprising five (5) animals per group. Group 1 served as the control and was not exposed to HS, while Groups 2, 3 and 4 were exposed to HS inside the heating chamber, regulated at $38\pm 1^\circ\text{C}$ for 2, 4 and 8 hours respectively for thirty 30 days. Animals were anaesthetized by cervical dislocation and blood was collected by cardiac puncture for haematological analysis: packed cell volume (PCV), haemoglobin concentration (Hb), red blood

*Corresponding author: E-mail: bruno.chinko@uniport.edu.ng;

cell (RBC), white blood cell (WBC), MID cells percentage (MID), lymphocytes, neutrophil and platelet counts were determined using a haematology auto-analyzer. Other haematological indices: mean corpuscular volume (MCV), mean corpuscular haemoglobin (MCH), mean corpuscular haemoglobin concentration (MCHC), neutrophil-lymphocyte ratio (NLR), platelet-lymphocyte ratio (PLR), total lymphocyte count (TLC), plateletcrit (PCT), mean platelet volume (MPV), platelet distribution width (PDW) and platelet large ratio (PCLR) were determined in line with standard formulae.

Results: Data from the study indicated a significantly raised PCV, RBC, Hb, NLR and PLR among the heat-exposed groups compared to the control ($p < 0.05$). Also, mean values of WBC, TLC, lymphocytes and MPV significantly decreased compared to the control ($p < 0.05$).

Conclusion: Long-term hemorheological changes are involved in the physiological adaptation to heat stress. According to the information from the current study, it appears that HS caused a rise in the PVC, Hb RBC, NLR, and PLR levels and a decrease in WBC, TLC, lymphocytes, and MPV. The data points to HS as a potential cause of increased blood viscosity, inflammation, and tissue damage, as well as immune system suppression and disruption of platelet synthesis and activation.

Keywords: Heat stress; haematology; red blood cell; white blood cell; mean platelet volume.

1. INTRODUCTION

In the past century, the earth's temperature has been gradually rising with strong evidence linking it to human activity, particularly the burning of fossil fuels such as coal, oil, and gas [1,2]. The impact of global warming and rising temperature on human physiology has continued to gain traction as the world continues to experience rising temperatures due to climate change [3,4]. Heat is a natural as well as an occupational hazard as it affects workers in various industries, particularly those who work outdoors or in hot environments [5,6]. With the human capacity to adapt to varied climates and environmental conditions in physiological and behavioural terms, there are clear and absolute limits to the amount of heat exposure an individual can tolerate [7,8].

Heat stress results when a change in body temperature exceeds the upper critical limits, producing a strain in the biological systems and overwhelming heat conservation/dissipation mechanisms, and affecting productivity and metabolic rate [9]. Homeothermic creatures regulate their body temperature within a narrow range and hence when heat is generated in the course of metabolic activity, a steady state is maintained by activating a heat loss mechanism to dissipate the excess heat [10,11]. Physiological adaptability to heat stress involves behavioural or metabolic modifications in response to heat to increase the dissipation of excess body heat to the environment to negate heat load in the body [12,13]. Thermoregulation involves a neural process that integrates external and internal information from the thermal

environment to an appropriate efferent response, allowing the organism to maintain a stable internal environment relative to a variable external environment [9,14,15]. The general homeostatic responses to HS stress include increased core body temperature, respiration rate, water consumption and peripheral vasoconstriction [16,17]. These responses are divided into two phases (acute and chronic) which correspond to the two stages of adaptation to heat stress. The acute response involves the activation of thermal receptors of the skin and hypothalamus which activates the autonomic nervous system leading to the release of glucocorticoids and catecholamines which modify metabolism. On the other hand, the chronic response is driven by continuous exposure to the HS causing changes in the homeorhetic and endocrine systems.

The adverse impact of heat has been fairly documented. As many intracellular molecular structures function on relatively weak interactions for stabilization, heat stress can cause significant cellular alterations and damage, impairing protein, DNA and RNA synthesis [18,19]. Also, heat stress increases oxidative stress by damaging the mitochondria and increasing intracellular reactive oxygen species (ROS) leading to protein and DNA damage as well increased lipid peroxidation [20-23]. Long-term exposure to heat stress can lead to endocrine responses involved in the release of stress hormones which can depress the thyroid hormones and hence affect energy utilization and lipid metabolism [24,25]. Furthermore, heat increases the level of plasma cortisol which down-regulates the activity of neutrophils,

depressing immunity [26]. Despite the various studies on the impact of heat stress, there is a paucity of data regarding the effect of heat stress on haematological parameters. Available data show varying and conflicting results. The present study, therefore, aims to evaluate the effect of heat stress on haematological profile using heat-stressed Wistar rat models.

2. MATERIALS AND METHODS

2.1 Research Animals

Twenty (20) apparently healthy male Wistar rats (200-250g, 12-16 weeks) were sourced from the animal house of the Department of Human Physiology, University of Port Harcourt and used for the study. The animals were allowed to acclimatize for two (2) weeks before the start of the study. The animals were housed under standard animal husbandry conditions in well-ventilated, clean wooden cages with optimal conditions: 12 hr day/night cycle, temperature 28 - 30°C, humidity 45 – 50%. The animals had access to standard rat chow and water *ad libitum*.

2.2 Heat Chamber

Heat stress (HS) was simulated using a perforated heated wooden chamber (30cm x 50cm x 25cm). The chamber was heated using a non-light heat emitter ceramic bulb (Simple Duluth xe, China), fitted with a digital thermometer (Shenzhen Brav Electronic Technologies Co., Ltd, China) and regulated using a heat switch (Popu Electric, China). The chamber was maintained at 38±1°C.

2.3 Research Design and Heating Protocol

Twenty (20) male Wistar rats were weighed and randomly divided into four (4) groups of five (5) animals each. During the experiment, Group 1 animals were used as the control and were kept inside a heating chamber without access to food or water, but they were not exposed to HS. Groups 2, 3 and 4 served as the study groups and were exposed to HS inside the heating chamber with the temperature regulated at 38±1°C for 2, 4 and 8 hours respectively. After each heating session, the animals were removed from the heating chamber and allowed free access to rat chow and water while they cooled passively at an ambient temperature of 26 – 30°C. Daily controlled heat exposure lasted for thirty (30) days.

2.4 Laboratory Analysis

Animals were anaesthetized by cervical dislocation and blood was collected by cardiac puncture and transferred into an EDTA sample bottle for haematological assay. The packed cell volume (PCV), haemoglobin concentration (Hb), red blood cell (RBC), white blood cell (WBC), MID cells percentage (MID), lymphocytes, neutrophil and platelet counts were determined using a haematology auto-analyzer (Automatic Haematology Analyzer, Mindray, China). Other haematological indices: mean corpuscular volume (MCV), mean corpuscular haemoglobin (MCH), mean corpuscular haemoglobin concentration (MCHC), neutrophil-lymphocyte ratio (NLR), platelet-lymphocyte ratio (PLR), total lymphocyte count (TLC), plateletcrit (PCT), mean platelet volume (MPV), platelet distribution width (PDW) and platelet large ratio (PCLR) were calculated in line with standard formulae [27-31].

The animals utilized in the study were housed and managed according to standard guidelines for the ethical and humane treatment of laboratory animals during experimental use [30,31].

2.5 Statistical Analysis

Data obtained from laboratory investigations were analyzed using IBM Statistical Product and Service Solutions (SPSS version 25). The mean and standard error of the mean of each parameter was calculated for each research group. The mean values obtained for study groups (2,3 & 4) were compared to the control (Group 1) using the analysis of variance (ANOVA) followed by a least significant difference (LSD) post hoc analysis (ANOVA). A p-value less than 0.05 ($p < 0.05$) was considered statistically significant.

3. RESULTS

The effects of heat stress on the PVC, Hb, RBC, MCV and MCHC of male Wistar rats are shown in Table 1. There was a significant increase in the PCV, Hb and RBC among the HS animals compared to the control ($p < 0.05$).

Table 2 shows the effect of HS on the WBC, lymphocyte, neutrophil, MID cell percentage, NLR, PLR and TLC of male Wistar rats. There was a significant reduction in WBC, lymphocytes, and TLC among the HS rats compared to the

control ($p < 0.05$). Conversely, the HS rats had a significantly increased NLR and PLR compared to the control ($p < 0.05$).

The effect of HS on PLT and other platelet indices is shown in Table 3. Although the PLT, PCT, PDW and PCLR all showed a marginal decrease among the HS animals when compared to the control ($p > 0.05$), only the MPV showed a significant reduction for 4 hr HS animals compared to the control ($p < 0.05$).

4. DISCUSSION

Heat stress occurs due to rising global temperatures and the exposure of certain

industrial workers to hot ambient temperatures [9,14]. Physiological adaptability to heat stress involves long-term metabolic and hemorheological modifications [9,12,15]. The present study evaluated the effect of a 30 days-controlled HS on the haematological profile of Wistar rats. Haematological parameters serve as vital indicators of the overall health status and the functional state of the body. They also serve as physiological gauges to changes in environmental temperature, stress and diseases [32,33]. The plasma fluid serves as a thermoregulatory medium to reduce hyperthermia with alterations in erythrocytic, leucocytic and immunologic variables [26,34].

Table 1. Effect of heat stress on packed cell volume, haemoglobin concentration, red blood cell count and red cell indices of male Wistar rats

Parameters	Control (n=5)	2 hr HS (n=5)	4 hr HS (n=5)	8 hr HS (n=5)
Packed cell volume (%)	44.30±0.63	49.66*±1.68	48.32*±1.26	51.08*±1.11
Haemoglobin concentration (g/dl)	12.94±0.21	13.68±0.16	13.92*±0.29	14.46*±0.33
Red blood cell ($\times 10^{12}/L$)	7.05±0.09	8.38*±0.77	7.67±0.18	7.86±0.10
Mean corpuscular volume (fL)	62.81±0.80	60.35±3.27	62.10±0.64	64.93±0.64
Mean corpuscular haemoglobin (pg)	18.34±0.23	16.80±1.32	18.16±0.26	18.38±0.19
Mean corpuscular haemoglobin concentration (g/dL)	29.22±0.36	27.69±1.05	28.83±0.22	28.31±0.17

Result is given as mean \pm standard error of mean; *significantly different compared to control ($p < 0.05$)

Table 2. Effect of heat stress on total white blood cell count, white blood cell differential count and some haematological indices of male Wistar rats

Parameters	Control (n=5)	2 hr HS (n=5)	4 hr HS (n=5)	8 hr HS (n=5)
White blood cells ($\times 10^9/L$)	4.00±0.64	1.62*±0.27	4.12±0.80	4.04±0.98
Lymphocytes ($\times 10^9/L$)	3.20±0.50	1.56*±0.42	2.58±0.45	3.22±0.73
Neutrophil ($\times 10^9/L$)	0.36±0.10	0.22±0.02	0.28±0.07	0.52±0.12
MID cells percentage ($\times 10^9/L$)	0.24±0.07	0.10±0.03	0.20±0.04	0.30±0.10
Neutrophil-Lymphocyte ratio	2.7±0.58	12.15*±4.26	4.16±0.86	4.88±1.51
Platelet-Lymphocyte ratio	232.53±36.95	573.02*±173.04	277.33±49.60	245.93±44.01
Total Lymphocyte Count	339.59±51.36	134.28*±25.29	351.68±76.57	322.77±72.77

Result is given as mean \pm standard error of mean; *significantly different compared to control ($p < 0.05$)

Table 3. Effect of Heat Stress on platelet count and other platelet parameters of male Wistar rats

Parameters	Control (n=5)	2 hr HS (n=5)	4 hr HS (n=5)	8 hr HS (n=5)
Platelets ($\times 10^9/L$)	676.80±35.17	668.80±55.49	636.80±47.26	667.20±48.57
Plateletcrit (%)	0.58±0.24	0.74±0.24	0.51±0.04	0.54±0.04
Mean platelet volume (fL)	8.66±0.20	8.34±0.25	8.06*±0.21	8.14±0.06
Platelet distribution width (%)	12.32±0.75	11.78±0.98	10.54±0.66	11.04±0.26
Platelet large ratio	19.86±1.91	17.60±2.18	15.48±1.59	15.68±0.53

Result is given as mean \pm standard error of mean; *significantly different compared to control ($p < 0.05$)

Data from the present study showed that animals exposed to HS had significantly raised PCV, Hb and RBC compared to the non-heat exposed control ($p < 0.05$) (Table 1). The packed cell volume (PCV) is a measure of the proportion of blood cells (majorly RBCs) in whole blood relative to the plasma. Heat stress can cause dehydration which shrinks the size of the plasma, thereby elevating the PCV [35]. Furthermore, when exposed to high temperatures, the body may experience heat-induced polycythemia, a condition where the number of red blood cells (RBCs) increases as a response to heat stress [36]. Since RBCs are responsible for transporting oxygen to the body's tissues, the body may produce more RBCs to meet the elevated demand caused by heat. Also, exposure to HS leads to the activation of the sympathetic nervous system and the subsequent release of catecholamines such as epinephrine and norepinephrine. These hormones trigger the contraction of the spleen, resulting in the release of stored red blood cells into the bloodstream. As a result, there is an increase in packed cell volume (PVC), red blood cell count, and haemoglobin concentration [35,37]. It has also been hypothesized that the increase in PVC and Hb in response to elevated high temperature could be due to an increase in the availability of nutrients required for haemoglobin synthesis as the animals may have consumed more food due to HS [35,38]. This increase in PCV, RBC and Hb is similar to earlier findings [33,39,40]. However, Srikanthakumar et al. and Sivakumar et al. [41,42] observed a decrease in PCV RBC and Hb concentration in heat-stressed goats and sheep respectively.

The study observed a significantly reduced WBC, TLC and lymphocytes among the 2 hours HS animals compared to the non-exposed control group ($p < 0.05$) (Table 2). WBCs are very significant components of the immune system which help the body to fight against infections and other foreign bodies [32,43]. Exposure to HS may have suppressed either the production of white blood cells (WBCs) in the bone marrow or their survival in circulation. This can lead to a reduction in the number of available WBCs to combat infections [40]. Furthermore, HS can lead to the activation of the hypothalamic-pituitary-adrenal (HPA) axis, which leads to the release of cortisol. Cortisol is a stress hormone that can suppress the immune system, leading to a decrease in the number of lymphocytes [26,33,34,44]. Data from the study also indicate a significantly increased neutrophil-lymphocyte

ratio (NLR) and platelet-lymphocyte ratio (PLR) among HS animals compared to the control ($p < 0.05$). NLR is employed as an easy and reliable marker of immune response to various infectious and non-infectious stimuli as it incorporates the innate immune response mainly due to neutrophils and adaptive immunity supported by lymphocytes [45,46]. Elevated NLR suggests systemic inflammatory response (SIRS) due to bacterial infection, cancer, stroke, severe trauma and tissue damage [47]. HS has been shown to cause tissue damage characterized by elevated serum malondialdehyde (MDA) [13,48]. The platelet-lymphocyte ratio (PLR) is a novel inflammatory biomarker used in the prediction of inflammation and mortality with higher values suggesting systemic inflammation, atherosclerosis and platelet activation [30,49]. The present study hypothesizes that HS may have depressed immunity and increased systemic inflammation.

According to the study, the platelet counts and other platelet indices exhibited a slight decrease, but the reduction was not statistically significant. There was a significant reduction in the mean platelet volume (MPV) in animals subjected to 4 hours of HS compared to the control group ($p < 0.05$). Mean platelet volume (MPV) is a measure of the average size of platelets in the blood. Platelets play a crucial role in blood clotting and inflammation. Dehydration and changes in blood viscosity can affect platelet size and function [50]. The MPV have been investigated in connection with inflammation and platelet activation [51,52]. Also, HS can trigger SIRS with inflammatory mediators such as interleukin-6 (IL-6) and C-reactive protein (CRP) changing the size and shape of platelets, promoting platelet destruction or even interfering with platelet production [53,54]. This suggests that HS may be associated with increased SIRS and platelet dysfunction.

5. CONCLUSION

Physiological adaptability to heat stress involves long-term hemorheological modifications. Based on the available data from the present study, it appears that HS led to elevated levels of PVC, RBC, NLR, and PLR and a reduction in WBC, TLC, lymphocytes and MPV. The evidence suggests that HS could be responsible for increasing blood viscosity, inflammation and tissue damage, depressing immune function and disrupting the production and activation of platelets.

CONSENT

It is not applicable.

ETHICAL APPROVAL

The research protocol and design were approved by the University of Port Harcourt Research Ethics committee.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Soeder DJ, Soeder DJ. Fossil fuels and climate change. Fracking and the Environment: A scientific assessment of the environmental risks from hydraulic fracturing and fossil fuels. 2021:155-85.
2. Letcher TM. Why do we have global warming? Managing Global Warming: Elsevier. 2019:3-15.
3. Amoo LM, Fagbenle RL. Climate change in developing nations of the world. Applications of Heat, Mass and Fluid Boundary Layers: Elsevier. 2020:437-71.
4. Dincer I. Renewable energy and sustainable development: A crucial review. Renewable and Sustainable Energy Reviews. 2000;4(2):157-75.
5. Spector JT, Masuda YJ, Wolff NH, Calkins M, Seixas N. Heat exposure and occupational injuries: review of the literature and implications. Current environmental health reports. 2019;6:286-96.
6. Ioannou LG, Foster J, Morris NB, Piil JF, Havenith G, Mekjavic IB, et al. Occupational heat strain in outdoor workers: A comprehensive review and meta-analysis. Temperature. 2022;9(1):67-102.
7. Kovats RS, Hajat S. Heat stress and public health: A critical review. Annu. Rev. Public Health. 2008;29:41-55.
8. Addo-Bediako A, Chown SL, Gaston KJ. Thermal tolerance, climatic variability and latitude. Proceedings of the Royal Society of London. Series B: Biological Sciences. 2000;267(1445):739-45.
9. Collier RJ, Baumgard LH, Zimbelman RB, Xiao Y. Heat stress: Physiology of acclimation and adaptation. Animal Frontiers. 2019;9(1):12-9.
10. Cheung SS, McLellan TM, Tenaglia S. The thermophysiology of uncompensable heat stress. Sports Medicine. 2000;29(5):329-59.
11. Gonzalez-Rivas PA, Chauhan SS, Ha M, Fegan N, Dunshea FR, Warner RD. Effects of heat stress on animal physiology, metabolism, and meat quality: A review. Meat Science. 2020;162:108025.
12. Rashamol VP, Sejian V, Bagath M, Krishnan G, Archana PR, Bhatta R. Physiological adaptability of livestock to heat stress: An updated review. Journal of Animal Behaviour and Biometeorology. 2020;6(3):62-71.
13. Chinko BC, Umeh OU. Alterations in lipid profile and oxidative stress markers following heat stress on wistar rats: Ameliorating role of vitamin C. Biomedical Sciences. 2023;9(1):12-7.
14. Nakamura K, Morrison SF. A thermosensory pathway that controls body temperature. Nature Neuroscience. 2008;11(1):62-71.
15. Seebacher F, Franklin CE. Physiological mechanisms of thermoregulation in reptiles: A review. Journal of Comparative Physiology B. 2005;175:533-41.
16. Mortola JP, Frappell PB. Ventilatory responses to changes in temperature in mammals and other vertebrates. Annual Review of Physiology. 2000;62(1):847-74.
17. Sarangi S. Adaptability of goats to heat stress: A review. The Pharma Innovation Journal. 2018;7(4):1114-26.
18. Caspani M, Savioli M, Crotti S, Bruzzone P, Gattinoni L. Heat stress: Characteristics, pathophysiology and avoidable mistakes. Minerva Anestesiologica. 2004;70(7-8): 617-24.
19. Roti Roti JL. Cellular responses to hyperthermia (40–46 C): Cell killing and molecular events. International Journal of Hyperthermia. 2008;24(1):3-15.
20. Halliwell B, Cross CE. Oxygen-derived species: Their relation to human disease and environmental stress. Environmental Health Perspectives. 1994;102(suppl 10): 5-12.
21. Storey KB. Oxidative stress: Animal adaptations in nature. Brazilian Journal of Medical and Biological Research. 1996;29:1715-33.
22. Lin H, Du R, Zhang Z. Peroxide status in tissues of heat-stressed broilers. Asian-Australasian Journal of Animal Sciences. 2000;13(10):1373-6.

23. Yanay O, Gilad E. Heat injury. *Pediatric critical care*: Elsevier; 2011:1472-6.
24. Mantha L, Palacios E, Deshaies Y. Modulation of triglyceride metabolism by glucocorticoids in diet-induced obesity. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*. 1999;277(2):R455-R64.
25. Wen X, Wu W, Fang W, Tang S, Xin H, Xie J, et al. Effects of long-term heat exposure on cholesterol metabolism and immune responses in growing pigs. *Livestock Science*. 2019;230:103857.
26. Habeeb A, El-Tarabany A, Gad A, Atta M. Negative effects of heat stress on physiological and immunity responses of farm animals. *Change*. 2018;16(2.0):0-2.
27. Bain BJ, Bates I, Laffan MA. *Dacie and lewis practical haematology e-book*: Elsevier Health Sciences; 2016.
28. Rozenberg G. *Microscopic haematology: A practical guide for the laboratory*. Elsevier Australia; 2011.
29. Faria SS, Fernandes Jr PC, Silva MJB, Lima VC, Fontes W, Freitas-Junior R, et al. The neutrophil-to-lymphocyte ratio: A narrative review. *Ecancermedicalscience*. 2016;10.
30. Balta S, Ozturk C. The platelet-lymphocyte ratio: A simple, inexpensive and rapid prognostic marker for cardiovascular events. *Platelets*. 2015;26(7):680-1.
31. Kwon H-C, Kim SH, Oh SY, Lee S, Lee JH, Choi H-J, et al. Clinical significance of preoperative neutrophil-lymphocyte versus platelet-lymphocyte ratio in patients with operable colorectal cancer. *Biomarkers*. 2012;17(3):216-22.
32. Chinko BC, Pughikumo DT. Haematological and hepatorenal alterations induced by potash (akanwu) on male wistar rats. *International Blood Research & Reviews*. 2023;14(1):38-46.
33. Ocheja OB, Ayo JO, Aluwong T, Minka NS. Ameliorative effects of L-glutamine on haematological parameters in heat-stressed red sokoto goats. *Journal of Thermal Biology*. 2020;90:102571.
34. Okoruwa MI. Effect of heat stress on thermoregulatory, live bodyweight and physiological responses of dwarf goats in southern Nigeria. *European Scientific Journal*. 2014;10(27):255-64.
35. Al-Dawood A. Towards heat stress management in small ruminants-a review. *Annals of Animal Science*. 2017;17(1):59-88.
36. Sawka MN, Gonzalez RR, Young AJ, Muza SR, Pandolf KB, Latzka WA, et al. Polycythemia and hydration: Effects on thermoregulation and blood volume during exercise-heat stress. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*. 1988;255(3):R456-R63.
37. Stewart IB, McKenzie DC. The human spleen during physiological stress. *Sports Medicine*. 2002;32:361-9.
38. Rana MS, Hashem MA, Sakib MN, Kumar A. Effect of heat stress on blood parameters in indigenous sheep. *Journal of the Bangladesh Agricultural University*. 2014;12(1):91-4.
39. Alam MM, Hashem MA, Rahman MM, Hossain MM, Haque MR, Sobhan Z, et al. Effect of heat stress on behavior, physiological and blood parameters of goat. *Progressive Agriculture*. 2011;22(1-2):37-45.
40. Odo RI, Onoja SO, Osuagwu CO. Impact of heat stress on blood and serum biochemistry parameters in rats. *Notulae Scientia Biologicae*. 2019;11(3):347.
41. Srikanthakumar A, Johnson EH, Mahgoub O. Effect of heat stress on respiratory rate, rectal temperature and blood chemistry in Omani and Australian Merino sheep. *Small Ruminant Research*. 2003;49(2):193-8.
42. Sivakumar AVN, Singh G, Varshney VP. Antioxidants supplementation on acid base balance during heat stress in goats. *Asian-Australasian Journal of Animal Sciences*. 2010;23(11):1462-8.
43. Chinko BC, Amah-Tariah FS, Ekenna IC. Evaluation of the effects of calabash chalk on the haematological profile of Wistar rats. *Notulae Scientia Biologicae*. 2022;14(3):11281.
44. Inbaraj S, Sejian V, Bagath M, Bhatta R. Impact of heat stress on immune responses of livestock: A review. *Pertanika Journal of Tropical Agricultural Science*. 2016;39(4).
45. Zahorec R. Neutrophil-to-lymphocyte ratio, past, present and future perspectives. *Bratislavske Lekarske Listy*. 2021;122(7):474-88.
46. Song M, Graubard BI, Rabkin CS, Engels EA. Neutrophil-to-lymphocyte ratio and mortality in the United States general population. *Scientific Reports*. 2021;11(1):464.
47. Buonacera A, Stancanelli B, Colaci M, Malatino L. Neutrophil to lymphocyte ratio:

- An emerging marker of the relationships between the immune system and diseases. *International Journal of Molecular Sciences*. 2022;23(7): 3636.
48. Malyar RM, Li H, Liu D, Abdulrahim Y, Farid RA, Gan F, et al. Selenium/Zinc-Enriched probiotics improve serum enzyme activity, antioxidant ability, inflammatory factors and related gene expression of wistar rats inflated under heat stress. *Life Sciences*. 2020;248: 117464.
 49. Gunaldi M, Goksu S, Erdem D, Gunduz S, Okuturlar Y, Tiken E, et al. Prognostic impact of platelet/lymphocyte and neutrophil/lymphocyte ratios in patients with gastric cancer: A multicenter study. *International Journal of Clinical and Experimental Medicine*. 2015; 8(4):5937.
 50. Sloop GD, De Mast Q, Pop G, Weidman JJ, Cyr JAS. The role of blood viscosity in infectious diseases. *Cureus*. 2020;12(2).
 51. Yuri Gasparyan A, Ayvazyan L, Mikhailidis DP, Kitas GD. Mean platelet volume: a link between thrombosis and inflammation? *Current Pharmaceutical Design*. 2011;17(1):47-58.
 52. Park Y, Schoene N, Harris W. Mean platelet volume as an indicator of platelet activation: Methodological issues. *Platelets*. 2002;13(5-6):301-6.
 53. Schmoeller D, Picarelli MM, Paz Munhoz T, Poli de Figueiredo CE, Staub HL. Mean platelet volume and immature platelet fraction in autoimmune disorders. *Frontiers in Medicine*. 2017;4:146.
 54. Lippi G, Franchini M. Platelets and immunity: the interplay of mean platelet volume in health and disease. *Taylor & Francis*. 2015:555-7.

© 2023 Umeh and Chinko; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/97945>