



# Nitrogen Uptake and Use Efficiency of Aerobic Rice as Affected by Different Nitrogen Management: A Review

Vaishali Singh <sup>a++</sup>, Chandrakant Chaubey <sup>a++\*</sup>, Anil Kumar <sup>b#</sup>,  
Narinder Panotra <sup>ct</sup>, Apurba Mishra <sup>d</sup>, Dipankar Barman <sup>e</sup>,  
Himanshu Sekhar Behera <sup>f++</sup>  
and Sharan S P <sup>g‡</sup>

<sup>a</sup> Department of Soil Science, Sardar Vallabhbhai Patel University of Agriculture & Technology, Meerut, India.

<sup>b</sup> P.G. Department of Chemistry, Sahibganj College Sahibganj -816109, Jharkhand, India.

<sup>c</sup> Institute of Biotechnology, SKUAST Jammu, J&K, India.

<sup>d</sup> Faculty of Agricultural Sciences, SOADU, Bhubaneswar, India.

<sup>e</sup> Indian Institute of Soybean Research, Indore, India.

<sup>f</sup> Department of Soil Science and Agricultural Chemistry, Lovely Professional University, Punjab-144411, India.

<sup>g</sup> Division of Agricultural Physics, ICAR-Indian Agricultural Research Institute, Pusa, New Delhi-110012, India.

## Authors' contributions

The sole author designed, analysed, interpreted and prepared the manuscript.

## Article Information

DOI: <https://doi.org/10.9734/jeai/2024/v46i92803>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/121530>

Review Article

Received: 11/06/2024

Accepted: 13/08/2024

Published: 24/08/2024

<sup>++</sup> Research Scholar;

<sup>#</sup> Head of Department;

<sup>†</sup> Associate Professor;

<sup>‡</sup> Ph. D Research Scholar;

\*Corresponding author: E-mail: [chandrakantchaubey1997@gmail.com](mailto:chandrakantchaubey1997@gmail.com)

**Cite as:** Singh, Vaishali, Chandrakant Chaubey, Anil Kumar, Narinder Panotra, Apurba Mishra, Dipankar Barman, Himanshu Sekhar Behera, and Sharan S P. 2024. "Nitrogen Uptake and Use Efficiency of Aerobic Rice As Affected by Different Nitrogen Management: A Review". *Journal of Experimental Agriculture International* 46 (9):45-60. <https://doi.org/10.9734/jeai/2024/v46i92803>.

## ABSTRACT

A significant amount of the world's population relies heavily on rice, which is grown in both lowland and upland rice systems. Irrigated lowland rice systems account for over 76% of the world's rice production. Although the aerobic rice system can be viewed as a revolutionary substitute for conventional rice, increased nitrogen (N) losses and decreased N-use efficiency (NUE) have made it difficult for this novel method to be widely adopted. One of the most crucial ingredients in the cultivation of rice is nitrogen (N). Less than 50% of N is recovered in both lowland and highland systems. Soil erosion, volatilisation, leaching, and denitrification account for the majority of the loss of applied nitrogen. Consequently, it will be prudent to maximise the use of nitrogenous fertilisers in order to achieve high yields and maintain a level of production costs that can be sustained. For rice cultivation in the aforementioned scenario, nitrogenous fertiliser needs to be used more wisely and efficiently. This review focusses on the various production practices that can increase the efficiency with which N is used, such as liming acid soils, providing N at appropriate rates, using appropriate sources, applying N at the right time and method, rotating crops, using cover crops, implementing conservation tillage, planting N-efficient genotypes.

*Keywords: Aerobic rice; rice ecosystems; nutrient use efficiency; nitrogen management; agronomic practices.*

## 1. INTRODUCTION

“In order to fulfil the future food demands of the expanding global population, rice (*Oryza sativa* L.) is regarded as a staple food for over half of the world's population. However, maintaining productivity amid resource constraint is difficult. Although rice is grown and consumed on every continent, the majority of it is grown and consumed in Asia, namely in China and India, which are the two largest producers and consumers of rice” [1]. “In addition, natives and immigrants from Asia, Africa, and South America consume enormous amounts of rice in North America and Europe. Studies have shown that in order to meet the demands of an expected rapidly growing population due to economic and industrial developments, as well as rapid population growth, global rice production will need to increase by approximately 20% by 2030 and 30% by 2050, at a rate of 0.6–1% increase annually” [2]. “For several years, agronomists have prioritised their study on examining different approaches to enhance rice grain production” [3,4]. “Given that the aerobic rice system may be viewed as a transformative substitute for conventional rice, increased nitrogen (N) losses and decreased N-use efficiency (NUE) have made it difficult for this novel method to be widely adopted” [5]. Increased N levels are necessary for agricultural plants to go through regular growth and developmental cycles.

“N is a mineral nutrient that is necessary for healthy plant development and metabolism since it is a major component of amino acids, nucleic acids, and many photosynthetic metabolites”

[6,7]. “However, uncontrolled and erratic N fertiliser application has increased the environmental sustainability concerns associated with higher N losses through volatilisation or leaching, causing pollution in the environment due to greenhouse effects and accelerating the depletion of the ozone layer” [8,9]. “Although aerobic rice growth and yield are enhanced by high N fertiliser application, environmental and economic sustainability are jeopardised. A number of environmental problems associated with increased nitrogen losses as nitrous oxide (N<sub>2</sub>O), ammonia (NH<sub>3</sub>), and nitrate (NO<sub>3</sub><sup>-</sup>) have been brought to light by the irregular and uncontrolled use of N fertilisers” [10]. This poses a threat to environmental sustainability because of the increased potential for global warming, ozone depletion, and eutrophication of water resources. Therefore, improving NUE in aerobic rice has emerged as a critical requirement for the creation of a long-term production system and predicts the economic return, is an important soil–plant interaction feature that has been studied for a long time.

Agronomic management approaches that are appropriate are necessary to achieve the efficient utilisation of synthetic N fertilisers, as irregular agricultural management practices contribute significantly to global anthropogenic emissions [11,12]. This review examines the assessment of several agricultural and agronomic management strategies designed to lower elevated nitrogen losses while increasing the NUE and grain production in an aerobic rice system with minimal risks to the environment.

Here are some key ways aerobic rice can help rice grow with less water: The following are some of the ways through which aerobic rice can assist rice to grow with the least water:

**Water savings:** aerobic rice fields are 50% less water consumptive than flood-borne paddy fields. Leakage involves a significant quantity of water as some of this water is lost through seepage, percolation and evaporation. Aerobic soil minimizes such losses of water to a great extent.

**Effective root development:** Aerobic rice lacks flooded and anaerobic conditions hence it can develop long, strong and branched roots that possess the potential to grow throughout the soil profile in search of water and nutrients. This is because the major branch roots thicken and grow deeper into the ground thus making the plant less susceptible to water stress.

**Higher water efficiency:** Aerobic rice has a higher efficiency in water uptake compared to flooded rice through root and stomatal pathways. This helps the stored water in the soil to be usable for longer durations between the irrigation exercises.

**Soil structure retention:** This is because the nature of waterlogged soils is that they wash away the structure of the soil with time. The wet and dry conditions that aerobic rice subject the soils to, help to retain a good structural and porosities that enable water retention and availability.

**Rainfall:** This is also capable of infiltrating and being stored in the soil if the soil type is appropriate. This rainfall can then be used by aerobic rice, whereas in flooded paddies, 70–90% of the rainfall infiltrates into the water and is unavailable for plant uptake.

In general, aerobic rice is more adaptable to the use of available soil moisture through root system and plant physiological processes to transpire less water than flooded rice and the plant utilizes rainfall for their production to reach high yields with significantly lower water compared to the flooded rice.


## 2. ADOPTING AEROBIC RICE SYSTEM

About 30% of the freshwater on Earth is used by the flooded rice system, and over 45% of it is used in Asia [13]. Due to the accelerated

demand for water from various sectors, freshwater resources are becoming more scarce, endangering the sustainability of conventional rice and making it more vulnerable to the effects of climate change. This necessitates the development of novel water-saving technologies through various modifications without compromising public health or environmental safety [14]. Traditionally, the rice-growing technique involves submerged circumstances, with standing water that is between 5 and 10 cm deep over the whole rice-growing season. Due to a lack of resources, including labour, electricity, and irrigation water, traditional rice farming is becoming more and more problematic.

For typical rice systems, puddling is an essential technique. However, it degrades the soil's structure, making it more difficult to prepare the land for subsequent crops that require more energy to achieve optimal soil tilth [15]. Consequently, these variables highlight how urgently traditional systems need to be modified through the introduction of water-saving techniques, including growing aerobic rice, which may function as an eco-efficient system by conserving natural resources without compromising grain output (Fig. 1).

Since rice naturally reacts negatively to drought, a decrease in irrigation supplies may result in a production fall [17]. Raised bed cultivation [13], the alternate wetting and drying system (AWD) [18], saturated soil culture [19], and the system of rice intensification (SRI) [20] are some of the modified technologies that have been developed in previous studies to reduce water application in rice systems. Due to the ongoing needs for puddling and soil saturation during crop growth, some of these are not water-use efficient and cannot receive a significant amount of water input. Rice is cultivated using the water-saving method of aerobic rice cultivation, which involves soil that is neither saturated nor puddled [16]. In addition to being adaptable in highland locations when additional irrigational water supplies are guaranteed, this modified method primarily targets irrigated lowland areas where irrigational water is insufficient for rice agriculture [21]. Early studies on aerobic rice technology reported grain yields of up to 6.5 t ha<sup>-1</sup> with water savings of about 60% [22]. These results showed that aerobic rice cultivation could be a large-scale, sustainable method of producing rice in the face of finite natural resources and climate change.



**Fig. 1. Benefits of transformation of conventional rice to aerobic rice system**

Source adapted from Joshi et al. [16]

### 3. NITROGEN (N) TRANSFORMATION IN RICE FIELD

Rice plants in paddy fields receive fertilisation from N-containing substances mostly in the form of urea. The hydrolysis of urea produces ammonium ( $\text{NH}_4^+$ ), which is then converted to  $\text{NO}_3^-$  in the oxic soil that surrounds the roots of the rice plants. Next,  $\text{NO}_3^-$  was incorporated into the oxidation of methane, denitrification, and anaerobic  $\text{NH}_4^+$  oxidation (anammox) after diffusing to the underlying anoxic soil [23]. As soil particle surfaces are usually negatively charged, cations (such as  $\text{NH}_4^+$ ) can be electrostatically attracted to them. Oxidising  $\text{NH}_4^+$  to hydroxylamine ( $\text{NH}_2\text{OH}$ ), nitrite ( $\text{NO}_2^-$ ), and finally  $\text{NO}_3^-$  is the process of nitrification. Two classes of chemoautotrophic bacteria, namely those that oxidise ammonia and nitrite (AOB and NOB), catalyse these activities. The rate-limiting step is the oxidation of  $\text{NH}_4^+$  to  $\text{NH}_2\text{OH}$ , which is catalysed by ammonia monooxygenase (AMO). From there,  $\text{NH}_2\text{OH}$  is oxidised to  $\text{NO}_2^-$  and subsequently to  $\text{NO}_3^-$  by nitrite oxidoreductase and hydroxylamine oxidoreductase, in that order [24]. Denitrification is the opposite process, wherein bacteria, fungi, and archaea decrease

$\text{NO}_3^-$  to  $\text{NO}_2^-$ , nitric oxide (NO),  $\text{N}_2\text{O}$ , and  $\text{N}_2$ . The environment will suffer from the partial decrease of  $\text{NO}_3^-$  to  $\text{N}_2\text{O}$  since  $\text{N}_2\text{O}$  is a greenhouse gas and has a global warming coefficient 300 times greater per molecule than  $\text{CO}_2$ , the primary ozone-destroying agent. Anammox, in which  $\text{N}_2$  is released from  $\text{NO}_2^-$  and  $\text{NH}_3$  by bacteria via the intermediates NO and hydrazine ( $\text{N}_2\text{H}_4$ ), respectively, is another reduction of  $\text{NO}_3^-$  to  $\text{NH}_3$  via  $\text{NO}_2^-$  under anaerobic or low-oxygen conditions known as dissimilatory nitrate reduction to ammonia (DNRA) [25].


According to Schimel and Bennett [26], nitrogen mineralisation is the process by which organic matter is converted by microbial activities into inorganic molecules (firstly,  $\text{NH}_4^+$ ). The rate of mineralisation is constrained by the depolymerisation of organic N, which releases organic N that is biologically available, such as amino acids. Microorganisms and plants can both absorb amino acids [27], however plants can only absorb most amino acid N after it has been converted to inorganic N [26]. In addition to nitrogen (N) taken by crop plants, a significant amount of N is retained in soil particles via mineralisation, which can release N for use in

subsequent crop cycles, and is immobilised by soil microbes, resulting in N storage in soil organic matter pools [28]. There are several variables impacting oxidation/reduction processes in the complicated N cycle in rice fields [24] (Fig. 2). The amount, variety, and makeup of microbial communities—such as those made up of bacteria, archaea, and fungi—as well as their interactions with roots might influence how nitrogen is transformed in rice paddies [29]. According to Zhong et al. [30], the metagenomic analysis revealed that the soil from the high-yielding rice field has a higher abundance of genes involved in the nitrification process and more taxa with N metabolism functions. This promotes the effective transformation of  $\text{NH}_4^+$  to  $\text{NO}_3^-$  in rice fields and stimulates high expression of  $\text{NO}_3^-$  transporters in plant roots. Soil microbiology is essential to the transformation of N in rice paddies.

#### 4. NITROGEN (N) LOSSES FROM RICE FIELD

Deprotonation of  $\text{NH}_4^+$  and volatilisation of  $\text{NH}_3$  gas to the atmosphere account for the

majority of nitrogen loss from paddy rice fields (globally, 10%–50% of applied fertilizer–N). A comparable amount (5.6%–50%) is lost through nitrification of N, which is followed by leaching and runoff because anion  $\text{NO}_3^-$  is poorly bound to soil. Denitrification (i.e.,  $\text{N}_2$ ,  $\text{N}_2\text{O}$ , and  $\text{NO}_x$  to the ambient air) accounts for a relatively smaller portion (0.03%–0.68%) of nitrogen loss [31]. In soils and solutions, ammonia and  $\text{NH}_4^+$  are in pH-dependent equilibrium, and volatilisation happens when there is an abundance of  $\text{NH}_4^+$  in the soil and a high soil alkalinity [32]. Additionally, plants can release gaseous fluxes of  $\text{NH}_3$ , which are most common in plants with high N contents and mostly occur in rice between anthesis and maturity [32]. With very few exceptions,  $\text{NO}_3^-$  does not bind to soil after  $\text{NH}_4^+$  transition to  $\text{NO}_3^-$ . As an anion,  $\text{NO}_3^-$  readily percolates through agricultural soil profiles and into groundwater. In soil,  $\text{NO}_3^-$  has a diffusion capacity that is 10–100 times greater than  $\text{NH}_4^+$  [33]. Air pollution issues are exacerbated by the production and degradation of  $\text{NO}_3^-$  by bacteria and archaea, which also releases additional  $\text{NO}_x$  and  $\text{N}_2\text{O}$ , potent greenhouse gases.



**Fig. 2. Nitrogen (N) cycling in rice production system**  
Source adapted from Coskun et al. [24]

## 5. NITROGEN (N) UPTAKE IN AEROBIC RICE ECOSYSTEM


The amount of N absorbed is measured as concentration, which is the amount of N in one unit of dry matter. N concentration in plants is often expressed as  $\text{g kg}^{-1}$  or as a percentage [34]. Concentration values are typically used to diagnosis plants for nutritional excess, shortage, or sufficiency. Nutrient concentrations from various agroecological zones can be extrapolated or utilised to diagnose nutritional problems in the same crop species. This is feasible because every element influencing nutrient availability depends on plant nutrient absorption. A measure of nutrient absorption is obtained by multiplying dry matter or grain yield by concentration; the result is given in accumulation or uptake units. For macronutrients, the unit of nutrient intake or accumulation in the field is  $\text{kg ha}^{-1}$ , whereas for micronutrients, it is  $\text{g ha}^{-1}$ . Nutrient absorption values are correlated with crop production levels and serve as helpful markers of the depletion of soil fertility. Dry matter accumulation was followed by nutrient accumulation patterns in agricultural plants, including rice [35]. Further nitrogen uptake and transformation are describing below.

Rice has the highest rate of N uptake, excluding K. The two inorganic types of nitrogen that rice predominantly absorbs are  $\text{NO}_3^-$  and  $\text{NH}_4^+$ . Since the environment is anaerobic,  $\text{NH}_4^+$  is typically thought of as the primary N type. Nevertheless, rice absorbs around 40% of the total N in the form of  $\text{NO}_3^-$  due to rhizosphere nitrification [36]. The absorption of nitrate has been well examined, and the associated rice transporters, such as the >80 nitrate transporter/peptide transporter (NRT1/PTR), 4 NRT2, and 2 nitrate assimilation related 2 (NAR2) members, have all been well-characterized. The first  $\text{NO}_3^-$  transporter with low affinity is OsNPF8.9 (OsNRT1) [37]. With the exception of NRT1.1, a dual affinity transporter, the majority of NRT1s exhibit poor affinity for  $\text{NO}_3^-$  [38]. High-affinity  $\text{NO}_3^-$  transporter genes were found in four NRT2 and two NAR2 genes. It was shown that nitrate transporters were effective in raising NUE. For example, up-regulated expression of OsNRT2.3b boosted grain output and NUE above wild type by 40% and enhanced the efficiency in  $\text{NO}_3^-$ ,  $\text{NH}_4^+$ , P, and Fe absorption [39]. While  $\text{NH}_4^+$  has long been thought to be the primary N absorption mechanism for rice plants, not much progress has been made in understanding  $\text{NH}_4^+$  transporters [40].

According to Li et al. [40] (2017), rice possesses at least 12 putative  $\text{NH}_4^+$  transporters (AMTs). Ragel et al. [41] suggest that overexpressing OsAMT1;1 might enhance  $\text{NH}_4^+$ , particularly in situations where  $\text{NH}_4^+$  levels are inadequate. The simultaneous activation of Glutamate synthetase 1 (OsGOGAT1) and OsAMT1/2 boosted the plant's ability to withstand N restriction and improved N remobilization and  $\text{NH}_4^+$  absorption overall [42]. Nevertheless, in contrast to the advancements in  $\text{NO}_3^-$  transporter research, the effectiveness of the  $\text{NH}_4^+$  transporter has not been as successfully improved [42]. After rice plants ingest  $\text{NO}_3^-$ , nitrate reductase (NR) in the cytoplasm converts it to  $\text{NO}_2^-$  first. Next, nitrite reductase (NiR) transfers  $\text{NO}_2^-$  to the plasmids, where it is further reduced to  $\text{NH}_4^+$  (Fig. 3). In the GS-GOGAT cycle, glutamine synthetase (GS) and GOGAT transform inorganic  $\text{NH}_4^+$  into glutamine. There are two types of GOGAT, ferredoxin-dependent (Fd-GOGAT, OsFd-GOGAT) and NADH-dependent (NADH-GOGAT, OsNADH-GOGAT1, and OsNADHGOGAT2) enzymes with varying electron donor specificity, and three GS isoforms (OsGS1;1, OsGS1;2, and OsGS1;3) [43].

## 6. NITROGEN USE EFFICIENCY (NUE) IN AEROBIC RICE ECOSYSTEM

The complicated feature of NUE is influenced by several chemical and metabolic routes as well as environmental factors [44]. Various agricultural main research wings have varied definitions of NUE. For example, plant physiologists consider numerous processes in the management of nitrogen, such as absorption, assimilation, allocation, and remobilization of nitrogen, which is particularly evident during leaf senescence [22]. Additionally, they split the NUE according to the several N cycle components, including N intake, N assimilation, N allocation, and N remobilization [45]. Grain yield equivalent to the entire quantity of accessible N from soil, including N fertiliser treatment, is what agronomists refer to as the NUE [46]. All agricultural experts worldwide concur that NUE is made up of two essential components that are readily evaluated in both controlled and field settings: the N uptake efficiency and the N utilisation efficiency [47]. N utilisation efficiency is the native ability of the plant to use the absorbed N, facilitating absorption and remobilization to create end harvest products. On the other hand, N uptake efficiency is the total capability to absorb, or uptake, N given from soil N pools.



**Fig. 3. Schematic illustrating the nitrate and ammonium assimilation and metabolism in plants. Source adapted from Li et al. [40]**

The best way to characterise N absorption efficiency is as the best possible balance between N remobilization and assimilation efficiencies [47]. In the meantime, N remobilization efficiency is determined by the total amount of N that is remobilised between the source and sink tissues, whereas N assimilation efficiency is defined as the overall capacity for the assimilation of inorganic N to manufacture amino acids and various other essential N-containing molecules [48]. According to Ierna and Mauromicale [49], each component of NUE is associated with a variety of complicated features, such as root morphological parameters, leaf senescence, N remobilization, and the ability to take accessible N from soil N pools. Enhanced N absorption and remobilization, enhanced N intake by various genetic alterations, and their control are the key drivers that can improve NUE. Furthermore, to raise NUE, a mix of agronomic techniques and traditional breeding methods is essential.

### 6.1 Important Elements for Improving Nitrogen Use Efficiency (NUE)

To meet the future food needs of the anticipated population growth, improvements in NUE under

aerobic rice systems and other non-N<sub>2</sub> fixing crops are needed. These improvements can be made through other crops, irrigation, and fertiliser management techniques, as well as improved and modified N-use efficient varieties [50]. The use of controlled-release N fertilisers, urease inhibitors for synthetic N fertilisers, crop variety management in terms of water and NUE, plant N uptake management, irrigation techniques, fertiliser input management, and other agronomic integrative crop management approaches are some of the management practices that are focused on improving plant N uptake and NUE [51]. Research on NUE has shown that it is primarily related to the kind and application of fertiliser inputs, irrigation techniques, soil N availability, and plant N absorption [52]. The linked management strategy, which combines irrigation and nitrogen control, is a widely used and environmentally sustainable technology in contemporary agriculture that may maintain rice output in the face of limited water resources while maintaining environmental sustainability. When N and irrigational managements are combined, N losses and fertiliser application rates are decreased, and NUE and irrigational water inputs are improved by over 20% [53]. Consequently, raising the total NUE in aerobic

rice systems requires a thorough comprehension and assessment of all the aforementioned parameters influencing NUE and their management.

## **7. AGRICULTURAL NITROGEN (N) MANAGEMENT FOR AEROBIC RICE ECOSYSTEM**

“Since the amount of N harvested in grains and soil N pools varies between intensive and extensive approaches, agricultural management aiming to increase NUE without compromising crop productivity should be in accordance with local soil and climatic conditions. This could involve intensive and extensive soil and crop residue management practices” [54]. Tillage practices present another difficulty for aerobic rice systems. While successive no-tillage or regular tillage methods led to higher N- and water-use efficiencies, they also increased crop competition because of increased weed invasion [55]. Therefore, for increased NUE and crop production in aerobic rice systems, integrated crop management strategies together with sophisticated irrigation control techniques are needed. In the section that follows, a few agricultural management techniques to raise NUE are covered.

### **7.1 Site-Specific Nutrient Management**

“A strategy known as “site-specific nutrient management” (SSNM) was developed in recent decades with the goal of producing fertiliser recommendations that are particular to a given field and improving the nutrient-use efficiency of important nutrients like phosphorus (P), potassium (K), and nitrogen (N)” [56]. “In order to synchronise nutrient provision and demand based on differences in plant requirements, indigenous nutrient provision, and nutrient recovery from applied fertilisers and other resources, the SSNM is a dynamic, typically plant-based, soil and crop growth season-specific nutrient management approach. Increasing nutrient-use efficiency is the main goal of SSNM in order to boost crop yields and local farmers’ economic outputs” [57]. For smallholder rice farmers in Asia, where field occupations are typically small and there are significant spatial variations in soil nutrient status and management, SSNM was first developed. This method of managing nutrients is primarily predicated on the idea of estimating nutrient requirements through the comparison of the total amount of nutrients needed to achieve a given

crop’s production target and the native supply of that particular nutrient [58]. Therefore, by adjusting the time of N fertiliser application to meet the plant’s prospective N requirements, this technique might promote the optimal use of N fertilisers in aerobic rice systems, enhance NUE, and prevent environmental concerns [57].

### **7.2 Integrated Nutrient Management Strategies**

To improve N recovery and NUE, integrated nutrient management (INM) entails making the best and most balanced use of naturally occurring N components, such as plant wastes, organic manures, biological N fixation, mineral N fertilisers, and their complementing interacting pathways [59]. The enhancement of the physico-chemical properties of the soil [60] or the improvement of plant root growth and the supply of necessary N along with other micronutrients [61] are the two main causes of the beneficial effects of the INM approach, which involves the mixed use of organic (such as biochar, manures, and compost) and inorganic N resources. To fully capitalise on the beneficial effects that are essential to raising net economic returns to farmers in terms of targeted production while preserving soil quality and raising NUE, it is imperative that the interacting pathways under INM be adequately evaluated and understood. A notable rise in crop yield and NUE may result from the complementing interactions between N from synthetic and non-synthetic fertilisers and other micronutrients. Therefore, improved NUE, reduced environmental hazards, and higher aerobic rice production will result from the wise and balanced use of N fertilisers from all available resources (organic and inorganic) through INM approaches. This will help ensure rice sustainability in the future even in the face of resource scarcity due to climate change.

### **7.3 Incorporation of Enhanced Use Efficiency N Fertilizers**

As of right now, a number of fertiliser products are useful because they may increase the NUE of applied N fertilisers by reducing N losses associated with the rice production system. The main characteristics of these fertiliser products are their ability to either slow down the rate at which N is released or to interfere with N transformation pathways in order to minimise N losses. N fertilisers with regulated and slow release have a crucial role in mitigating different types of nitrogen losses, which in turn affects



recovery and availability of nitrogen [62]. NO<sub>3</sub>-containing N fertilisers are more prone to leaching than NH<sub>4</sub><sup>+</sup> containing fertilisers, whereas the former is more vulnerable to volatilisation loss. A variety of commercially available slow- and controlled-release nitrogen fertilisers can be used in conjunction with possible mitigation techniques to improve NUE and decrease N losses, ultimately leading to a sustainable agricultural production system [63]. Because of their ability to postpone N release patterns, these slow- and controlled-release N compounds can enhance the synchronisation between soil N supply and plant N requirements, hence reducing N losses. Furthermore, commercial urea coated with neem can be extensively employed in aerobic rice systems because to its proven slow-release N fertiliser properties in South Asia [64]. Measures that can increase the use of these compounds are necessary because of the higher manufacturing costs and restricted availability of such controlled-release N fertilisers, which make their broad use difficult.

#### 7.4 Residue Management

The primary source and sink for the C and N cycles are crop wastes. By first converting nitrogen into inorganic forms and then mineralising it at later phases of crop growth when crop need for nitrogen becomes significant, the incorporation of crop wastes into the soil assures that the plants will have access to nitrogen for a longer length of time. Furthermore, because soil microorganisms temporarily immobilise nutrients released into the soil and preserve them in slowly accessible forms, including crop leftovers into the aerobic rice system may be advantageous [54]. It has previously been shown that plant residues from a variety of cereals can guarantee the provision of N between 40 and 100 kg ha<sup>-1</sup> season<sup>-1</sup> and help the soil system accumulate organic C, which enhances the NUE and soil N pool even more [65]. Legumes have a high N content and lower C/N ratios than cereals, therefore adding them to the aerobic rice system can be a useful way to increase soil N pools [66]. Furthermore, half of the total quantity of N would be accessible for plant absorption after two months of residue integration due to the faster rates of mineralisation of legume plant residues [67,68].

#### 7.5 Green Manuring

Numerous legume crops possess an exceptional ability to fix atmospheric nitrogen, making them

suitable for use in green manuring within aerobic rice systems [69]. According to Sharma et al. [70], in order for legume crops to be considered superior for green manuring, they need to have certain key characteristics. These include the ability to produce large amounts of dry matter quickly and easily, the potential to fix atmospheric free nitrogen, the ability to grow quickly and shorter growth durations for easy settlement into an intensive cropping system, and the cultivation of minimal land management practices. However, the quality and amount of residues available, soil characteristics, microbial diversity and activity, soil moisture content, and atmospheric factors all have a major role in the positive effects and effectiveness of legumes as green manures. To improve N recovery and total NUE, it is therefore essential to cultivate crops in a recurrent succession on the same plot of land and to properly rotate legumes with aerobic rice systems. It has been discovered that legume crops have the ability to lessen the requirement for subsequent crops in the rotation system. Additionally, by increasing total carbon stocks, which in turn increases plant availability of N, narrow C:N ratios of legume residues can improve the physical, chemical, and biological properties of the soil [71].

#### 7.6 Precision Farming

In order to maximise net profits, crop production, resource efficiency, and system sustainability in terms of quality and quantity, precision farming is an information- and technology-based farm input management system that focusses on the use of cutting-edge technologies and principles in identifying, analysing, and managing spatial and temporal variabilities associated with all characteristics of crop production systems [72-74]. According to Yousefi and Razdari [75], one important way to increase the NUE under varying climatic conditions is to measure variabilities in the field with respect to the availability of N for plant uptake and apply the optimal dose of N fertilisers at the right time by using technologies like rate applicator, remote sensing, geographic information system (GIS), and global positioning system (GPS). In order to assess a crop's need for more N, modern crop management systems can also make use of local or distant N sensors. Consequently, increasing the NUE in intensive agricultural systems may benefit from the application of the aforementioned technology and precision agricultural techniques.

## 8. CLIMATE CHANGE: CONTEXT AND CHALLENGES IN AEROBIC RICE PRODUCTION

There is evidence of climate change, which has an effect on ecosystems, the population, and livelihoods [76]. Climate change poses significant developmental issues for the global society due to its extensive effects, especially for the impoverished and people in developing and underdeveloped nations that depend on natural resources [77]. Communities are affected by the global phenomena of climate change in different ways within and across nations [78]. In a nation, the impoverished, politically disenfranchised and marginalised communities, and those with the least resources are the most likely to be negatively impacted by climate change and are also the least likely to diversify their sources of income [79]. "The mean earth surface temperature has been rising since the start of the 19th century due to anomalies and unevenness in the variabilities of climatic components" [80]. "The data from the Intergovernmental Panel on Climate Change (IPCC) annual report 6 (AR6) working group I (WGI) report show that, based on projected temperature fluctuations over the last few decades, the global atmospheric temperature has risen by 1.09°C since the 1950s. The majority of this temperature change was observed in the last 30 years of the 19th century" [81]. There are three main causes that are known to contribute to climate change: land-use changes, human-induced factors like greenhouse gas emissions (GHGs), and natural fluctuations.

Assessments of future climate change indicate that extreme weather events, temperature variations, and precipitation intensity, frequency, and patterns will all increase in severity in the years to come [82]. Concurrently, it has been predicted that increased oceanic absorption of CO<sub>2</sub> will cause the earth's surface temperature to rise more slowly than that indicated by climate model estimates [83]. Therefore, there are a number of errors in the patterns, frequency, and intensity of precipitation due to the climate models' inability to accurately estimate the hydrological cycle over the tropical areas [84]. One of the environmental factors that determines whether a crop plant will succeed or fail is the climate. It supports and is crucial to both short- and long-term crop planning, especially in the case of climatic disasters or altered weather patterns.

Uneven and unstable climate variations have put the world's rice crop in jeopardy, especially in Asia. Rice production and climate change are strongly related since the flooded rice system adds to greenhouse gas emissions and increases global warming, which in turn affects the rice agriculture system. The Food and Agriculture Organisation (FAO) forecasts that in order to achieve food security goals in the face of resource scarcity, rice production must rise by 40% by the end of the 2030s [85]. If this increase in food is achieved using the least amount of environmentally friendly practices, environmental pollution problems might get worse. One of the primary causes of greenhouse gas emissions is thought to be rice grown in traditional flooding settings [86]. In light of the impending shortage of resources and the need to reduce methane (CH<sub>4</sub>) emissions [87], a number of transformations have been implemented for the production of rice. However, these changes come with a number of environmental risks because of the substantial losses of nitrogen through volatilisation and leaching.

Generally speaking, the conventional rice system produces higher greenhouse gas emissions. However, when the conventional system is modified to meet food security goals and increase the sustainability of rice production in the face of limited natural resource availability, high nitrogen losses result from the intermixing of nitrification and denitrification processes [79]. Therefore, it is required to modify the traditional rice system or replace it with an aerobic rice system in order to reduce greenhouse gas emissions and meet future food requirements in the face of resource constraint [88]. However, widespread adaptation of aerobic rice necessitates research and evaluation of modifications in the N cycle pathways and microbial abundance and functioning under the aerobic rice system through agronomic management approaches due to high N losses and environmental pollution issues.

## 9. FUTURE THRUST

In order for the aerobic rice system to be widely adapted, study on the ways in which the interaction activities of microbial communities—such as nitrifiers, denitrifiers, and ammonia oxidizers—change under different soil conditions is required on a global scale. One of the main issues facing the aerobic rice system at the moment is our incomplete knowledge of the interactions between various microbial

populations when implementing N fertiliser control strategies. Furthermore, while implementing any agronomic management practices, there is still little ability to interactively evaluate changes in the abundance, diversity, and functioning of microbial communities in both space and time under aerobic soil conditions. Agronomic management practices for aerobic rice systems that incorporate NUE-boosting while N-reducing N losses may have an effect on the quantity and activity of soil microbial communities as well as biochemical characteristics. Therefore, more research on aerobic rice systems is urged, with an emphasis on designing experiments that evaluate changes in the variety and activity of microbial communities, plant N needs, and N management strategies in an interactive manner. This will increase our knowledge and comprehension of the ways that various agronomic N management techniques affect the quantity, diversity, and activities of soil microbial communities. This knowledge and comprehension may then be used to improve NUE and lower N losses for increased grain yield and environmental safety.

## 10. CONCLUSION

Crops depend on nitrogen, and adding it to aerobic rice production systems is crucial to boosting crop yield and keeping up with the growing human population. A significant amount of the nitrogen (N) that is applied to rice, however, is lost to the environment, which causes a number of issues. Although the absorption and transformation of nitrogen in the rice ecosystem is complicated and involves a wide range of entities and participants, there are several chances for N management intervention. In aerobic rice systems, providing the ideal conditions and making agronomic modifications to account for N losses can help close the production gap between traditional and conventional rice systems. When switching from an anaerobic to an aerobic rice system, there is a risk of substantial nitrogen losses because of changes in soil nitrogen pools, water availability, and microbial activity. In order to achieve widespread adoption of the aerobic rice system, rice cultivars that exhibit improved root characteristics for improved nitrogen uptake and utilisation, improved yield traits for lowland cultivars, and improved water stress tolerance traits for upland cultivars must be introduced. To sustain grain yield and ensure environmental sustainability in the face of future projected climate change and input scarcity, research-

based evaluation of agronomic management measures for aerobic rice systems is required to decrease N losses and increase NUE.

## DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

## COMPETING INTERESTS

Author has declared that no competing interests exist.

## REFERENCES

1. Peng S, Tang Q, Zou Y. Current status and challenges of rice production in China. *Plant Prod. Sci.* 2009;12:3–8. DOI: 10.1626/pps.12.3
2. Shew AM, Durand-Morat A, Putman B, Nalley LL, Ghosh A. Rice intensification in Bangladesh improves economic and environmental welfare. *Environ. Sci. Policy.* 2019;95:46–57. DOI: 10.1016/J.ENVSCI.2019.02.004
3. Mongiano G, Titone P, Pagnoncelli S, Sacco D, Tamborini L, Pilu R, et al. Phenotypic variability in Italian rice germplasm. *Eur. J. Agron.* 2020;120:126131. DOI: 10.1016/j.eja.2020.126131
4. Zhao H, Mo Z, Lin Q, Pan S, Duan M, Tian H, et al. Relationships between grain yield and agronomic traits of rice in southern China. *Chil. J. Agric. Res.* 2020;80:72–79. DOI: 10.4067/s0718-58392020000100072
5. Xu C, Yang F, Tang X, Lu B, Li Z, Liu Z, et al. Super rice with high sink activities has superior adaptability to low filling stage temperature. *Front. Plant Sci.* 2021;12:2111. DOI: 10.3389/FPLS.2021.729021/BIBTEX.
6. Balestrini R, Brunetti C, Chitarra W, Nerva L. Photosynthetic traits and nitrogen uptake in crops: Which is the role of Arbuscular Mycorrhizal fungi? *Plants.* 2020;9:1105. DOI: 10.3390/PLANTS9091105.
7. Sandhu N, Sethi M, Kumar A, Dang D, Singh J, Chhuneja P. Biochemical and genetic approaches improving nitrogen use efficiency in cereal crops: A Review. *Front. Plant Sci.* 2021;12:757.

- DOI: 10.3389/FPLS.2021.657629/BIBTEX.
8. Raza S, Zhou J, Aziz T, Afzal MR, Ahmed M, Javaid S, et al. Piling up reactive nitrogen and declining nitrogen use efficiency in Pakistan: A challenge not challenged (1961-2013). *Environ. Res. Lett.* 2018;13:034012.  
DOI: 10.1088/1748-9326/aaa9c5
  9. Devi OR, Laishram B, Singh S, Paul AK, Sarma HP, Bora SS and Devi SS. A Review on Mitigation of Greenhouse Gases by Agronomic Practices towards Sustainable Agriculture. *International Journal of Environment and Climate Change.* 2023;13(8):278–287.  
Available: <https://doi.org/10.9734/ijecc/2023/v13i81952>
  10. Fowler D, Coyle M, Skiba U, Sutton MA, Cape JN, Reis S, Sheppard LJ, Jenkins A, Grizzetti B, Galloway JN, Vitousek P, Leach A, Bouwman AF, Butterbach-Bahl K, Dentener F, Stevenson D, Amann M, Voss M. The global nitrogen cycle in the twenty-first century. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 2013;368:20130164.  
Available: <https://doi.org/10.1098/rstb.2013.0164>
  11. Smith LED, Siciliano G. A comprehensive review of constraints to improved management of fertilizers in China and mitigation of diffuse water pollution from agriculture. *Agric. Ecosyst. Environ.* 2015 ;209:15–25.  
DOI: 10.1016/J.AGEE.2015.02.016
  12. Aryal JP, Sapkota TB, Krupnik TJ, Rahut DB, Jat ML, Stirling CM. Factors affecting farmers' use of organic and inorganic fertilizers in South Asia. *Environ. Sci. Pollut. Res.* 2021;28:51480–51496.  
DOI:10.1007/S11356-021-13975-7/TABLES/10
  13. Kadiyala MDM, Mylavaram RS, Li YC, Reddy GB, Reddy MD. Impact of aerobic rice cultivation on growth, yield, and water productivity of rice-maize rotation in semiarid tropics. *Agron. J.* 2012;104:1757–1765.  
DOI: 10.2134/agronj2012.0148
  14. Rasul G. Managing the food, water, and energy nexus for achieving the Sustainable Development Goals in South Asia. *Environ. Dev.* 2016;18:14–25.  
DOI: 10.1016/J.ENVDEV.2015.12.001.
  15. Khairul Alam M, Bell RW, Hasanuzzaman M, Salahin N, Rashid MH, Akter N, et al. Rice (*Oryza sativa* L.) Establishment techniques and their implications for soil properties, global warming potential mitigation and crop yields. *Agronomy.* 2020;10:888.  
DOI: 10.3390/AGRONOMY10060888
  16. Joshi R, Singh B, Shukla A. Evaluation of elite rice genotypes for physiological and yield attributes under aerobic and irrigated conditions in Tarai areas of western Himalayan region. *Curr. Plant Biol.* 2018;13:45–52.  
DOI: 10.1016/J.CPB.2018.05.001
  17. Ahmadikhah A, Marufinia A. Effect of reduced plant height on drought tolerance in rice. *3 Biotech.* 2016;6:221.  
DOI:10.1007/S13205-016-0542-3/TABLES/7
  18. Howell KR, Shrestha P, Dodd IC. Alternate wetting and drying irrigation maintained rice yields despite half the irrigation volume, but is currently unlikely to be adopted by smallholder lowland rice farmers in Nepal. *Food Energy Secur.* 2015;4:144.  
DOI: 10.1002/FES3.58
  19. Singh B, Mishra S, Bisht S, Joshi R, Singh B, Mishra S, et al. Growing rice with less water: Improving productivity by decreasing water demand. *Rice Improv.* 2021;147–170.  
DOI: 10.1007/978-3-030-66530-2\_5
  20. Chapagain T, Riseman A, Yamaji E. Assessment of System of Rice Intensification (SRI) and Conventional Practices under Organic and Inorganic Management in Japan. *Rice Sci.* 2011;18:311–320.  
DOI: 10.1016/S1672-6308(12)60010-9
  21. Thakur AK, Mohanty RK, Singh R, Patil DU. Enhancing water and cropping productivity through Integrated System of Rice Intensification (ISRI) with aquaculture and horticulture under rainfed conditions. *Agric. Water Manag.* 2015;161:65–76.  
DOI: 10.1016/J.AGWAT.2015.07.008
  22. Sandhu N, Sethi M, Kumar A, Dang D, Singh J, Chhuneja P. Biochemical and Genetic Approaches Improving Nitrogen Use Efficiency in Cereal Crops: A Review. *Front. Plant Sci.* 2021;12:757.  
DOI: 10.3389/FPLS.2021.657629/BIBTEX.
  23. Kuypers MMM, Marchant HK, Kartal B. The microbial nitrogen-cycling network. *Nat. Rev. Microbiol.* 2018;16:263–276.  
Available: <https://doi.org/10.1038/nrmicro.2018.9>
  24. Coskun D, Britto DT, Shi W, Kronzucker HJ. How plant root exudates shape the

- nitrogen cycle. *Trends Plant Sci.* 2017b;22:661–673.  
Available:<https://doi.org/10.1016/j.tplants.2017.05.004>
25. Rütting T, Boeckx P, Müller C, Klemedtsson L. Assessment of the importance of dissimilatory nitrate reduction to ammonium for the terrestrial nitrogen cycle. *Biogeosciences.* 2011;8:1779–1791.  
Available:<https://doi.org/10.5194/bg-8-1779-2011>
  26. Schimel JP, Bennett J. Nitrogen mineralization: Challenges of a changing paradigm. *Ecology.* 2004;85:591–602.  
Available:<https://doi.org/10.1890/03-8002>
  27. Nasholm T, Ekblad A, Nordin A, Giesler R, Hogberg M, Hogberg P. Boreal forest plants take up organic nitrogen. *Nature.* 1998;392:914–916.  
Available:<https://doi.org/10.1038/31921>
  28. Johnson DW, Cheng W, Burke IC. Biotic and abiotic nitrogen retention in a variety of forest soils. *Soil Sci. Soc. Am. J.* 2000;64:1503–1514.  
Available:<https://doi.org/10.2136/sssaj2000.6441503x>
  29. Zhang H, Hou DP, Peng XL, Ma BJ, Shao SM, Jing WJ, Gu JF, Liu LJ, Wang ZQ, Liu YY, Yang JC. Optimizing integrative cultivation management improves grain quality while increasing yield and nitrogen use efficiency in rice. *J. Integr. Agric.* 2019a;18:2716–2731.  
Available:[https://doi.org/10.1016/S2095-3119\(19\)62836-4](https://doi.org/10.1016/S2095-3119(19)62836-4)
  30. Zhong YQW, Hu JH, Xia QM, Zhang SL, Li X, Pan XY, Zhao RP, Wang RW, Yan WM, Shanguan ZP, Hu FY, Yang CD, Wang W. Soil microbial mechanisms promoting ultrahigh rice yield. *Soil Biol. Biochem.* 2020;143:107741.  
Available:<https://doi.org/10.1016/j.soilbio.2020.107741>
  31. Coskun D, Britto DT, Shi W, Kronzucker HJ. Nitrogen transformations in modern agriculture and the role of biological nitrification inhibition. *Native Plants.* 2017a;3:1–10.  
Available:<https://doi.org/10.1038/nplants.2017.74>
  32. Schjoerring JK, Husted S, Mattsson M. Physiological parameters controlling plant-atmosphere ammonia exchange. *Atmos. Environ.* 1998;32:491–498.  
Available:[https://doi.org/10.1016/S1352-2310\(97\)00006-X](https://doi.org/10.1016/S1352-2310(97)00006-X)
  33. Robertson GP, Vitousek PM. Nitrogen in agriculture: Balancing the cost of an essential resource. *Annu. Rev. Environ. Resour.* 2009;34:97–125.  
Available:<https://doi.org/10.1146/annurev.environ.032108.105046>
  34. Fageria NK. Plant tissue test for determination of optimum concentration and uptake of nitrogen at different growth stages in lowland rice. *Commun Soil Sci Plant Anal.* 2003;34:259–270.
  35. Fageria NK. *Mineral nutrition of rice.* CRC Press, Boca Raton; 2014.
  36. Zhang Z, Chu C. Nitrogen-use divergence between indica and japonica rice: Variation at nitrate assimilation. *Mol. Plant.* 2019;13:6–7.  
Available:<https://doi.org/10.1016/j.molp.2019.11.011>
  37. Lin CM, Koh S, Stacey G, Yu SM, Lin TY, Tsay YF. Cloning and functional characterization of a constitutively expressed nitrate transporter gene, OsNRT1, from rice. *Plant Physiol.* 2000;122:379–388.  
Available:<https://doi.org/10.1104/pp.122.2.379>
  38. Gu JF, Li ZK, Mao YQ, Struik P, Zhang H, Liu LJ, Wang ZQ, Yang JC. Roles of nitrogen and cytokinin signals in root and shoot communications in maximizing of plant productivity and their agronomic applications. *Plant Sci.* 2018;274:320–331.  
Available:<https://doi.org/10.1016/j.plantsci.2018.06.010>
  39. Fan X, Tang Z, Tan Y, Zhang Y, Luo B, Yang M, Lian X, Shen Q, Miller AJ, Xu G. Overexpression of a pH-sensitive nitrate transporter in rice increases crop yields. *Proc. Natl. Acad. Sci. U.S.A.* 2016;113:7118–7123.  
Available:<https://doi.org/10.1073/pnas.1525184113>
  40. Li H, Hu B, Chu C. Nitrogen use efficiency in crops: Lessons from Arabidopsis and rice. *J. Exp. Bot.* 2017;68:2477–2488.  
Available:<https://doi.org/10.1093/jxb/erx101>
  41. Ragel P, Rodenas R, García-Martín E, Andres Z, Villalta I, Nieves-Cordones M, Rivero RM, Martínez V, Pardo JM, Quintero FJ, Rubio F. CIPK23 regulates HAK5-mediated high-affinity K<sup>+</sup> uptake in Arabidopsis roots. *Plant Physiol.* 2015;169:2863–2873.  
Available:<https://doi.org/10.1104/pp.15.01401>

42. Lee S, Marmagne, A., Park, J., Fabien, C., Yim, Y., Kim, S., Kim, T., Lim, P.O., MasclauxDaubresse, C., Nam, H.G., 2020. Concurrent activation of OsAMT1;2 and OsGOGAT1 in rice leads to enhanced nitrogen use efficiency under nitrogen limitation. *Plant J.* 103, 7–20. Available:https://doi.org/10.1111/tpj.14794.
43. Gu J, Yang J. Nitrogen (N) transformation in paddy rice field: Its effect on N uptake and relation to improved N management. *Crop and Environment.* 2022;1(1):7–14. Available:https://doi.org/10.1016/j.crope.2022.03.003
44. Farooq MS, Gyllbag A, Virk AL, Xu Y. Adaptability mechanisms of japonica rice based on the comparative temperature conditions of harbin and qiqihar, heilongjiang province of Northeast China. *Agronomy.* 2021;11:2367. DOI: 10.3390/agronomy11112367
45. Lee S. Recent advances on nitrogen use efficiency in rice. *Agronomy.* 2021;11:753. DOI: 10.3390/agronomy11040753.
46. The SV, Snyder R, Tegeder M. Targeting nitrogen metabolism and transport processes to improve plant nitrogen use efficiency. *Front. Plant Sci.* 2021;11:2330. DOI: 10.3389/FPLS.2020.628366/BIBTEX.
47. Zhao C, Ma G, Zhou L, Zhang S, Su L, Sun X, et al. Effects of nitrogen levels on gene expression and amino acid metabolism in Welsh onion. *BMC Genomics.* 2021;22:803. DOI: 10.1186/s12864-021-08130-y
48. Dellerio Y. Manipulating amino acid metabolism to improve crop nitrogen use efficiency for a sustainable agriculture. *Front. Plant Sci.* 2020;11:1857. DOI: 10.3389/FPLS.2020.602548/BIBTEX.
49. Ierna A, Mauromicale G. Sustainable and Profitable Nitrogen Fertilization Management of Potato. *Agronomy.* 2019;9:582. DOI: 10.3390/AGRONOMY9100582
50. Mboyerwa PA, Kibret K, Mtakwa PW, Aschalew A. Evaluation of growth, yield, and water productivity of paddy rice with water-saving irrigation and optimization of nitrogen fertilization. *Agronomy.* 2021;11:1629. DOI: 10.3390/agronomy11081629.
51. Ma X, Zhang F, Liu F, Guo G, Cheng T, Wang J, et al. An integrated nitrogen management strategy promotes open-field pepper yield, crop nitrogen uptake, and nitrogen use efficiency in southwest china. *Agriculture.* 2022;12:524. DOI: 10.3390/AGRICULTURE12040524
52. Rawal N, Pande KR, Shrestha R, Vista SP. Nutrient use efficiency (NUE) of wheat (*Triticum aestivum* L.) as affected by NPK fertilization. *Plos One.* 2022;17:e0262771. DOI: 10.1371/JOURNAL.PONE.0262771.
53. Yang Z, Li N, Ma P, Li Y, Zhang R, Song Q, et al. Improving nitrogen and water use efficiencies of hybrid rice through methodical nitrogen–water distribution management. *F. Crop. Res.* 2020;246:107698. DOI: 10.1016/J.FCR.2019.107698.
54. Sarkar S, Skalicky M, Hossain A, Brestic M, Saha S, Garai S, et al. Management of Crop Residues for Improving Input Use Efficiency and Agricultural Sustainability. *Sustainability.* 2020;12:9808. DOI: 10.3390/SU12239808
55. Habbib H, Hirel B, Spicher F, Dubois F, Tétu T. In Winter Wheat (*Triticum Aestivum* L.), No-Till Improves Photosynthetic Nitrogen and Water-Use Efficiency. *J. Crop Sci. Biotechnol.* 2019;231:39–46. DOI: 10.1007/S12892-019-0122-0
56. Chivenge P, Zingore S, Ezui KS, Njoroge S, Bunquin MA, Dobermann A, et al. Progress in research on site-specific nutrient management for smallholder farmers in sub-Saharan Africa. *F. Crop. Res.* 2022;281:108503. DOI: 10.1016/J.FCR.2022.108503
57. Rodriguez DGP. An assessment of the site-specific nutrient management (SSNM) strategy for irrigated rice in Asia. *Agriculture.* 2020;10:559. DOI: 10.3390/agriculture10110559.
58. Sharma S, Panneerselvam P, Castillo R, Manohar S, Raj R, Ravi V, et al. Web-based tool for calculating field-specific nutrient management for rice in India. *Nutr. Cycl. Agroecosyst.* 2019;113:21. DOI: 10.1007/S10705-018-9959-X
59. Sarwar N, Atique-ur-Rehman Farooq O, Wasaya A, Hussain M, El-Shehawi AM, et al. Integrated nitrogen management improves productivity and economic returns of wheat-maize cropping system. *J. King Saud Univ. Sci.* 2021;33:101475. DOI: 10.1016/J.JKSUS.2021.101475
60. Oladele S, Adeyemo A, Awodun M, Ajayi A, Fasina A. Effects of biochar and nitrogen fertilizer on soil physicochemical properties, nitrogen use efficiency and upland rice (*Oryza sativa*) yield grown on

- an Alfisol in Southwestern Nigeria. *Int. J. Recycl. Org. Waste Agric.* 2019;8:295–308. DOI:10.1007/S40093-019-0251-0/TABLES/6
61. Thind SS, Wu G, Tian M, Chen A. Significant enhancement in the photocatalytic activity of N, W co-doped TiO<sub>2</sub> nanomaterials for promising environmental applications. *Nanotechnology.* 2012;23:475706.
  62. Ransom CJ, Jolley VD, Blair TA, Sutton LE, Hopkins BG. Nitrogen release rates from slow- and controlled-release fertilizers influenced by placement and temperature. *Plos One.* 2020;15:e0234544. DOI: 10.1371/JOURNAL.PONE.0234544.
  63. Mahmud K, Panday D, Mergoum A, Missaoui A. Nitrogen losses and potential mitigation strategies for a sustainable agroecosystem. *Sustainability.* 2021;13:1–23. DOI: 10.3390/su13042400
  64. Naz MY, Sulaiman SA. Attributes of natural and synthetic materials pertaining to slow-release urea coating industry. *Rev. Chem. Eng.* 2017;33:293–308. DOI: 10.1515/REVCE-2015-0065/MACHINEREADABLECITATION/RIS
  65. Roozbeh M, Rajaie M. Effects of residue management and nitrogen fertilizer rates on accumulation of soil residual nitrate and wheat yield under no-tillage system in south-west of Iran. *Int. Soil Water Conserv. Res.* 2021;9:116–126. DOI: 10.1016/J.ISWCR.2020.09.007
  66. Moore KJ, Anex RP, Elobeid AE, Fei S, Flora CB, Goggi AS, et al. Regenerating Agricultural Landscapes with Perennial Groundcover for Intensive Crop Production. *Agronomy.* 2019;9:458. DOI: 10.3390/AGRONOMY9080458
  67. Wang L, Yuan X, Liu C, Li Z, Chen F, Li S, et al. Soil C and N dynamics and hydrological processes in a maize-wheat rotation field subjected to different tillage and straw management practices. *Agric. Ecosyst. Environ.* 2019;285:106616. DOI: 10.1016/J.AGEE.2019.106616.
  68. Devi OR, Ojha N, Laishram B, Dutta S, Kalita P. Roles of nano-fertilizers in sustainable agriculture and biosafety. *Environment and Ecology.* 2023;41(1B):457-463.
  69. Qaswar, M., Jing, H., Ahmed, W., Shujun, L., Dongchu, L., Lu, Z., et al. (2019). Substitution of Inorganic Nitrogen Fertilizer with Green Manure (GM) Increased Yield Stability by Improving C Input and Nitrogen Recovery Efficiency in Rice Based Cropping System. *Agronomy* 9:609. DOI: 10.3390/AGRONOMY9100609.
  70. Sharma S, Panneerselvam P, Castillo R, Manohar S, Raj R, Ravi V, et al. Web-based tool for calculating field-specific nutrient management for rice in India. *Nutr. Cycl. Agroecosyst.* 2019;113:21. DOI: 10.1007/S10705-018-9959-X.
  71. Vogel AM, Below FE. Hybrid selection and agronomic management to lessen the continuous corn yield penalty. *Agronomy.* 2018;8:228. DOI: 10.3390/agronomy8100228.
  72. van Evert FK, Gaitán-Cremaschi D, Fountas S, Kempenaar C. Can precision agriculture increase the profitability and sustainability of the production of potatoes and olives? *Sustainability.* 2017;9:1863. DOI: 10.3390/su9101863.
  73. Devi OR, Sarma A, Borah K, Prathibha RS, Tamuly G, Maniratnam K and Laishram B. Importance of zinc and molybdenum for sustainable pulse production in India. *Environment and Ecology.* 2023; 41(3C): 1853–1859. Available: <https://doi.org/10.60151/envec/lcch4556>.
  74. Laishram B, Singh TB, Kalpana A, Wangkheirakpam M, Chongtham SK and Singh W. Effect of Salicylic Acid and Potassium Nitrate on Growth and Yield of Lentil (*Lens culinaris L.*) under Rainfed Condition. *International Journal of Current Microbiology and Applied Sciences.* 2020;9(11):2779–2791. Available: <https://doi.org/10.20546/ijcmas.2020.911.337>
  75. Yousefi M, Razdari AM. Application of Gis and gps in Precision Agriculture (a Review). *Int. J. Adv. Biol. Biom. Res.* 2015;3:7–9.
  76. Stecula DA, Merkley E. Framing Climate Change: Economics, Ideology, and Uncertainty in American News Media Content from 1988 to 2014. *Front. Commun.* 2019;4:6. DOI:10.3389/FCOMM.2019.00006/BIBTEX
  77. Devkota N, Pajja N. Impact of climate change on paddy production: Evidence from Nepal. *Asian J. Agric. Dev.* 2020;17:63–78. DOI: 10.37801/AJAD2020.17.2.4.
  78. Ali S, Liu Y, Ishaq M, Shah T, Abdullah, Ilyas A et al. Climate change and its impact

- on the yield of major food crops: Evidence from Pakistan. *Foods*. 2017;6:39. DOI: 10.3390/FOODS6060039.
79. Islam SN, Winkel J. *Climate Change and Social Inequality*; 2017. Available: <http://www.ejnetindiaresource.org/ejissues/bali.pdf> (accessed on Apr 18, 2022).
80. Farooq MS, Wang X, Uzair M, Fatima H, Fiaz S, Maqbool Z, Rehman OU, Yousuf M, Khan MR. Recent trends in nitrogen cycle and eco-efficient nitrogen management strategies in aerobic rice system. *Frontiers in Plant Science*. 2022;13:960641.
81. IPCC. *Climate change 2021 the physical science basis summary for policymakers working group I contribution to the sixth assessment report of the intergovernmental panel on climate change*. Geneva: IPCC; 2021.
82. Tabari H. Climate change impact on flood and extreme precipitation increases with water availability. *Sci. Rep.* 2020;10:13768. DOI: 10.1038/s41598-020-70816-2.
83. Balmaseda MA, Hernandez F, Storto A, Palmer MD, Alves O, Shi L et al. The Ocean Reanalyses Intercomparison Project (ORA-IP). *J. Operat. Oceanogr.* 2015; 8:s80–s97. DOI: 10.1080/1755876X.2015.1022329.
84. Lorenz C, Kunstmann H, Lorenz C, Kunstmann H. The hydrological cycle in three state-of-the-art reanalyses: Intercomparison and performance analysis. *J. Hydrometeorol.* 2012;13:1397–1420. DOI: 10.1175/JHM-D-11-088.1
85. FAO. *OECD-FAO Agricultural Outlook 2017-2026 SpECiAl FOCuS: SOuthEASt ASiA*. Rome: FAO; 2016.
86. Oelbermann M. (ed.). *Sustainable agroecosystems in climate change mitigation*. Wageningen: Wageningen Academic Publishers; 2014. DOI: 10.3920/978-90-8686-788-2
87. Devi OR, Laishram B, Singh S, Paul AK, Sarma HP, Bora SS, Devi SS. A Review on Mitigation of Greenhouse Gases by Agronomic Practices towards Sustainable Agriculture. *International Journal of Environment and Climate Change*. 2023;13(8): 278–287. Available: <https://doi.org/10.9734/ijecc/2023/v13i81952>
88. Kiran Daggali G, Tiwari U, Pandey PK, Devi OR, Gireesha D, Laishram B, Patel A. Discovering new frontiers in plant breeding: The fascinating world of advancements shaping future growth. *International Journal of Research in Agronomy*. 2024;7(1):441–445. Available: <https://doi.org/10.33545/2618060x.2024.v7.i1f.262>

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/121530>